The first International African Symposium on Exotic Nuclei IASEN2013

Contribution ID: 59

Type: Oral

Investigating the photon strength function to discrete levels

Friday, 6 December 2013 11:45 (25 minutes)

Over the last decade several measurements in light- and medium-mass nuclei have reported an enhanced ability for the absorption and emission of gamma radiation (photon strength function PSF) at low energies. The impact of this effect may have profound implications on neutron capture reaction rates which are not only responsible for the formation of elements heavier than iron in stellar and supernova environments [1] but are also of central importance for advanced fuel cycles in nuclear reactors [2]. The results were received with significant skepticism by the community mainly due to the lack of any known mechanism responsible for such an effect but also because another established experimental technique failed to confirm the measurement. Now, a new experimental method which is free of model input and systematic uncertainties has been developed to determine the PSF. It is designed to study statistical feeding from the quasi-continuum (below the particle separation energies) to individual low-lying discrete levels. A key aspect to successfully study gamma decay from the region of high-level density is the detection and extraction of correlated high-resolution particle-gamma-gamma events which is accomplished using an array of Clover HPGe detectors and large area segmented silicon detectors. The excitation energy of the residual nucleus produced in the reaction is inferred from the detected proton energies in the silicon detectors. Gating on gamma-transitions originating from low-lying discrete levels specifies the states fed by statistical gamma-rays. Any particle-gamma-gamma event satisfying these and additional energy sum requirements ensures a clean and unambiguous determination of the initial and final states of the observed gamma rays. With these constraints the statistical feeding to discrete levels is extracted on an event-by-event basis.

In this talk I will review our experimental technique to extract information on the gamma-ray decay from the quasi-continuum and present results for 95Mo [3]. Furthermore, I will discuss on-going experimental efforts to explore the properties of statistical spectra at stable and radioactive beam facilities.

[1] M. Arnould, S. Goriely and K. Takahashi, Physics Reports 450, 97213 (2007).

[2] M.B. Chadwick et al., Data Nuclear Data Sheets 112, 2887 (2011).

[3] M. Wiedeking et al., Phys. Rev. Lett. 108, 162503 (2012).

Primary author: Dr WIEDEKING, Mathis (iThemba LABS)

Presenter: Dr WIEDEKING, Mathis (iThemba LABS)

Session Classification: Resonances Session