

4th China - South Africa Joint Symposium on Nuclear Physics



Report of Contributions

Contribution ID: 0

Type: **not specified**

Nuclear level densities and gamma-ray strength functions of $^{180,181,182}\text{Ta}$ and neutron capture cross sections

Most stable and extremely low abundance proton-rich nuclei with $A > 110$ are thought to be produced by the photodisintegration of s - and r -process seed nuclei. However, this so-called p -process is insufficient to explain the observed low abundance (0.012%) of the ^{180}Ta isotope. Hence combinations of several processes are considered to reproduce the observed abundance of ^{180}Ta in the cosmos, provoking debates and making it a unique case study. Significant uncertainties in the predicted reaction rates in p -nuclei arise due to large uncertainties in nuclear properties such as the nuclear level densities (NLD) and gamma-ray strength functions (gamma;SF) [1], as well as the actual astrophysical environments. An experiment was performed in October 2014 to extract the NLD and gamma;SF below the neutron threshold in $^{180,181,182}\text{Ta}$ isotopes which provide important input parameters for nuclear reaction models. In the present case study, these parameters were measured using the $^{181}\text{Ta}(^3\text{He}, ^3\text{He}')$ and $^{181}\text{Ta}(^3\text{He}, ^4\text{He})$ reactions with 34 MeV beam, $^{181}\text{Ta}(d, d')$ and $^{181}\text{Ta}(^3\text{He}, t)$ reactions with 15 MeV beam, and $^{181}\text{Ta}(d, d')$ and $^{181}\text{Ta}(d, p)$ reactions with 12.5 MeV beam at the Oslo Cyclotron Laboratory. Using the SiRi array at backward angles (64 silicon particle telescopes) and the CACTUS array (26 NaI(Tl) detectors), the NLD and gamma;SF were simultaneously extracted below the neutron separation energy from particle-gamma; coincidence matrices through iterative procedures using the Oslo method [2]. The experimental results have been used to determine the corresponding neutron capture cross sections, which in turn were utilized to extract Maxwellian averaged cross sections. The latter can be used in astrophysical network calculations to investigate the galactic production mechanism of ^{180}Ta . In this talk I will present results of this investigation of statistical properties for $^{180,181,182}\text{Ta}$ and the corresponding (n, gamma;) cross sections.

[1] S. Goriely et al., *A&A J.* 375, L35 (2001).

[2] A. Schiller et al., *Nucl. Instrum. Methods Phys. Res. A* 447, 498 (2000).

This work is based on the research supported in part by the National Research Foundation of South Africa Grant Number 92600 and the IAEA under research grant number 20454.

Primary author: Mr MALATJI, Kgashane (Physics Department, University of Stellenbosch, Matieland, 7602, South Africa)

Co-authors: Prof. GÖRGEN, Andreas (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); Dr LARSEN, Ann-Cecilie (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); Dr KHESWA, Bonginkosi (Department of Physics, University of Johannesburg, Johannesburg, 1850, South Africa); Mr BRITS, Christiaan (Physics Department, University of Stellenbosch, Matieland, 7602, South Africa); Dr BLEUEL, Darren (Lawrence Livermore National Laboratory, Livermore, California 94551, USA); Dr SAHIN, Eda (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); Mr ZEISER, Fabio (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); Dr GIACOPPO, Francesca (Helmholtz Institute Mainz, 55099 Mainz, Germany and GSI Helmholtzzentrum für Schwerionenforschung, 64291 Darmstadt, Germany); Mr BELLO GARROTE, Frank (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); Dr TVETEN, Gry (Department of Physics,

University of Oslo, NO-0316 Oslo, Norway); Dr NYHUS, Hilde (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); Dr HADYNSKA-KLEK, Kasia (INFN, Laboratori Nazionali di Legnaro Padova, 35020 Legnaro, Italy); Prof. GUTTORMSEN, Magne (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); Dr KLINTEFJORD, Malin (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); ROSE, Sunniva (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); Prof. SIEM, Sunniva (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); Dr RENSTRØM, Therese (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); Dr HAGEN, Trine (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); Mr INGEBERG, Vetle (Department of Physics, University of Oslo, NO-0316 Oslo, Norway); Dr WIEDEKING, mathis (Department of Subatomic Physics, iThemba LABS, Old Faure Road, 7131, South Africa)

Presenter: Mr MALATJI, Kgashane (Physics Department, University of Stellenbosch, Matieland, 7602, South Africa)

Contribution ID: 1

Type: **not specified**

Feasibility of Nuclear Plasma Interaction studies with the Activation Techniques

Electron-mediated nuclear plasma interactions (NPIs), such as Nuclear Excitation by Electron Capture (NEEC) or Transition (NEET), may have significant impact on nuclear cross sections in High Energy Density Plasmas (HEDPs). These HEDP environments are found in the cosmos where nucleosynthesis takes place. Attempts have failed so far in measuring the NEEC process [1], while NEET has recently been observed experimentally [2]. NEEC, NPIs have not been observed due to the narrowness of nuclear transitions ($\approx 1\text{eV}$). The NPIs may occur on highly excited nuclear states in the quasi-continuum which is populated in nuclear reactions prior to their decay by spontaneous γ -ray emission. Direct observation of NPIs are hindered by the lack of a clear signature of the effect in HEDP environments. Hence, a new signature [3] for NPIs on highly excited nuclei will be tested by investigating isomeric to ground state feeding from the quasi-continuum region. An experiment was performed using the reactions $^{197}\text{Au}(^{13}\text{C}, ^{12}\text{C})^{198}\text{Au}$ and $^{197}\text{Au}(^{13}\text{C}, ^{12}\text{C}2\text{n})^{196}\text{Au}$ at Lawrence Berkeley National Laboratory in inverse kinematics with a ^{197}Au beam of 8.5 MeV/u energy. The activated foils were counted at the low-background counting facility of Lawrence Livermore National Laboratory. I will discuss several measurements with different target configurations to investigate the feasibility of NPI studies.

Primary authors: Mr THEMBALETHU, Thembaletu (University of the Western Cape); Dr WIEDEKING, mathis (itl)

Presenter: Dr WIEDEKING, mathis (itl)

Contribution ID: 2

Type: **not specified**

PSF in $^{181,182}\text{Ta}$ and the emergence of the scissors resonance

Relatively small resonances on the low-energy tail of the giant electric dipole resonance such as the scissors or pygmy resonances can have significant impact on reaction rates. These rates are important input for modelling processes that take place in astrophysical environments and nuclear reactors. Recent results from the University of Oslo indicate the existence of a significant enhancement in the photon strength function for nuclei in the actinide region due to the scissors resonance [1]. Further, the M1 strength distribution of scissors resonances in rare earth nuclei has been studied extensively over the years [2]. In order to investigate the extent and persistence of the scissor resonance in other mass regions, an experiment was performed utilizing the NaI(Tl) gamma-ray detector array (CACTUS) and silicon particle telescopes (SiRi) at the cyclotron laboratory at the University of Oslo. Particle- γ coincidences from the $^{181}\text{Ta}(d,p)^{182}\text{Ta}$ reaction were used to measure the nuclear level density and photon strength function of the well-deformed ^{182}Ta system, to investigate the existence of resonances below the neutron separation energy. In this talk I will present and discuss the final results of this investigation and place our findings in the context of previous work.

[1] M. Guttormsen et al. Phys. Rev. Lett. 109, 162503 (2012).

[2] P. von-Neumann-Cosel, K. Heyde, and A. Richter, Rev. Mod. Phys., 82, 2365, (2010).

This work is based on the research supported in part by the National Research Foundation of South Africa Grant Number 92600.

Primary author: Mr BRITS, Christiaan (iThemba LABS)

Co-authors: Prof. GORGEN, A. (University of Oslo); Dr LARSEN, A.C. (University of Oslo); Mr KHESWA, Bonginkosi (iThemba LABS); Dr BLEUEL, D. (Lawrence Livermore National Laboratory, USA); Dr SAHIN, E. (University of Oslo); Mr ZEIZER, F.B. (University of Oslo); Mr BELLO GARROTE, F.L. (University of Oslo); Ms TVETEN, G.M. (University of Oslo); Dr F., Giacoppo (University of Oslo); Dr NYHUS, H.T. (University of Oslo); Dr HADYNSKA-KLEK, K (University of Oslo); Mr MALATJI, Kgashane (University of the Western Cape); Ms KLINTEFJORD, M (University of Oslo); Prof. GUTTORMSEN, M. (University of Oslo); Dr PAPKA, Paul (Stellenbosch University); Ms ROSE, S. (University of Oslo); Prof. SIEM, S. (University of Oslo); Ms RENSTROM, T. (University of Oslo); Ms HAGEN, T.W. (University of Oslo); Mr INGEBERG, V.W. (University of Oslo); Dr WIEDEKING, mathis (itl)

Presenter: Mr BRITS, Christiaan (iThemba LABS)

Contribution ID: 4

Type: **not specified**

SAIF - the South African Isotope Facility

The iThemba LABS' Radioactive-Ion Beam project has evolved into SAIF - the South African Isotope Facility. It has two phases, the first of which comprises the Low-Energy Radioactive-Ion Beam (LERIB) project and ACE Isotopes (Accelerator Centre for Exotic Isotopes) project. ACE isotopes calls for the installation of a commercial, off-the-shelf 70 MeV cyclotron for radionuclide production. It will remove isotopes production from the existing SSC accelerator, freeing additional beam time for research. The LERIB project is an upgraded version of the RIB "demonstrator", capable of producing neutron-rich beams of high-intensity, due to the fissioning of natural uranium at a rate of up to 6×10^{13} /s. The beams from LERIB will be of low-energy, 60 keV - suitable for decay studies and implantation in materials as radioactive probes. Phase 2 of the SAIF project is the Accelerator Centre for Exotic Beams (ACE Beams). It will see the addition of a post-accelerator, likely a LINAC, to take beams from the LERIB to high-energies for research into sub-atomic physics.

Primary author: Dr BARK, Robert (iThemba LABS)

Presenter: Dr BARK, Robert (iThemba LABS)

Contribution ID: 6

Type: **not specified**

TOWARDS IMPLEMENTATION/DEVELOPMENT OF STATE-OF-THE-ART ELECTRON SPECTROMETER CAPABILITY AT iThemba LABS

An electron spectrometer that implements a Si(Li) detector for the detection of internal conversion electrons and a combination of LaBr₃ detectors for the detection of coincident decays is undergoing development at iThemba LABS, South Africa. For optimization of measurements with the electron spectrometer, electron calibration sources (²⁰⁷Bi and ¹³³Ba) are used in the measurements of energy and momentum resolution of the Si(Li) detectors. Other parameters such as transmission and efficiency which are used in describing the performance of electron spectrometer [1], are also determined during the characterization measurements. Lifetime measurements which provide crucial information needed for the measurement of E0 matrix elements will be performed using LaBr₃ detectors. Electron energy versus magnet current calibration was done by varying the lens current repeatedly to ensure reproducibility. Characterization measurements are presented and discussed for the purpose of commissioning the electron spectrometer for the study of 0⁺ states at iThemba LABS.

Keywords: Electron spectrometer, internal conversion electrons, coincident decays, electron energy, magnetic current.

[1] J. Kantele, et al., Nuclear Instrument and Methods, 130(2): 467-474, 1975

Primary author: Mr AVAA, Abraham (iThemba/Wits)

Co-authors: Dr USMAN, Iyabo (Wits); Dr JONES, Pete (iThemba)

Presenter: Mr AVAA, Abraham (iThemba/Wits)

Contribution ID: 7

Type: **not specified**

Hybrid potential analysis of clustering in heavy and superheavy nuclei

Local core -cluster interaction of Saxon-Woods type functional form with parameters derived from Michigan-3-Yukawa (M3Y) microscopic potential model have been used to investigate the decay properties of heavy and superheavy nuclei.

Further applications of the potential model to the low lying spectra of heavy nuclei reveals its inadequacy in the internal structure consistent with earlier works.

The geometry of the interior is taken to be influenced by the possible core-cluster overlap.

An additional interaction in the interior taken to mimic the nucleon distribution in the overlap region is shown to improve the spectra.

Primary author: Dr IBRAHIM, taofiq (Stellenbosch University)

Co-author: Dr WYNGAARDT, Shaun (Stellenbosch University)

Presenter: Dr IBRAHIM, taofiq (Stellenbosch University)

Contribution ID: 8

Type: **not specified**

Pairing Isomers

The concept of configuration dependent pairing does not usually appear in the textbooks and is rarely considered in the literature. The resulting lowering of excited $0n+$ states into the pairing gap in even-even deformed nuclei accounts for many states that have traditionally been assumed to arise from time-dependent “ β vibrations” of the nuclear shape along the symmetry axis. The experimental evidence that requires the low-lying $0n+$ states to be re-evaluated will be reviewed from the early (p,t) work of Maher et al. [1] and Kolata and Oohoudt [2] to more recent γ -ray spectroscopy studies [3-5].

[1] J. V. Maher, J. R. Erskine, A. M. Friedman, J. P. Schiffer and R. H. Siemssen, Phys. Rev. Lett. 25, 302 (1970).

[2] J. J. Kolata and M. Oothoudt, Phys. Rev. 15, 1947 (1977).

[3] P. E. Garrett, J. Phys. G27, R1 (2001).

[4] W. D. Kulp et al., Phys. Rev. C77, 061301(R) (2008).

[5] J. F. Sharpey-Schafer et al., Eur. Phys. J. A47, 5 & 6 (2011).

Primary author: Prof. SHARPEY-SCHAFFER, John F (University of Western Cape)

Presenter: Prof. SHARPEY-SCHAFFER, John F (University of Western Cape)

Contribution ID: 9

Type: **not specified**

Searching for the low-energy enhancement in 91-Zr

The nuclear level density (NLD) and γ -ray strength function (SF) are quantities that give essential information about the behaviour of a nucleus at high excitation energy. NLD is denoted as the number of levels per unit of excitation energy. SF is denoted as a measure of the average reduced decay probability of a nucleus. These concepts are useful at high excitation energies where the spacing between the levels is small and gives information on degrees of freedom and underlying nuclear dynamics. The evidence of the low-energy enhancement in the SF for energies less than 4 MeV has been discovered in several nuclei. Recently, a strong enhancement of M1 transitions in ^{90}Zr has been predicted for γ -ray energies below 2 MeV in shell model calculations. In this work we explored the existence of the low-energy enhancement in the neighbouring ^{91}Zr isotope with the assumption that neighbouring isotopes have similar SF and hence provided first experimental NLD and SF for this nucleus. The experiment $^{90}\text{Zr}(d,p)^{91}\text{Zr}$ was conducted at the Oslo Cyclotron Laboratory (OCL). The SiRi (silicon telescope) array was used to detect charged ejectiles from the reaction. The CACTUS (NaI(Tl) detectors) array was utilized to detect rays that were in coincidence with charged particles. The nuclear level density and SF were extracted with the Oslo method. The existence of the LEE was observed, which agrees with the shell model calculations in ^{90}Zr . The NLD and SF quantities were used to calculate (n, γ) cross sections with the Talys reaction codes. These were compared with experimental data from direct measurement to test the reliability of the approach used in this work.

Primary author: Mr ZIKHALI, Bonginkosi Richard (University of Zululand)

Co-author: Dr WIEDEKING, mathis (itl)

Presenter: Mr ZIKHALI, Bonginkosi Richard (University of Zululand)

Contribution ID: 11

Type: **not specified**

Gamma-Ray Strength Function in ^{74}Ge from the Ratio Method

An increasing number of experiments reveal the presence of a low-energy enhancement in the gamma-ray strength function (GSF). The GSF, which is the ability of nuclei to absorb and emit γ rays, provides insight into the statistical properties of atomic nuclei. For this project the GSF was studied for ^{74}Ge which was populated in the reaction $^{74}\text{Ge}(p,p')^{74}\text{Ge}$ at a beam energy of 18 MeV. The data was collected with the STARS-LIBERACE array at Lawrence Berkeley National Laboratory. Silicon detector telescopes were used for particle identification and γ -rays in coincidence were detected with 5 Clover-type high-purity germanium detectors. Through the analysis particle- γ - γ coincidence events were constructed. These events, together with well-known energy levels, were used to identify primary γ -rays from the quasicontinuum. Primary γ -rays from a broad excitation energy region, which decay to two 0^+ states, six 2^+ states, two 3^+ states, five 3^- states, and four 4^+ states, could be identified. These states and the associated primary γ -rays are used to measure the GSF for ^{74}Ge with the Ratio Method [1], which entails taking ratios of efficiency corrected primary γ -ray intensities from the quasicontinuum. I will discuss the results from the analysis of the data from the above reaction and focus on the existence of the low-energy enhancement in ^{74}Ge . The results are further discussed in the context of other work done in ^{74}Ge using the (γ,γ') [2], $(^3\text{He},^3\text{He}')$ [3] and (α,α') [4] reactions.

Primary authors: Mr SOWAZI, Khanyisa (iThemba LABS); Dr WIEDEKING, mathis (itl)

Co-authors: Dr HURST, Aaron (Nuclear Science Division, Lawrence Berkeley National Laboratory); Dr GORGEN, Andreas (Department of Physics, University of Oslo); Dr GOLDBLUM, Bethany (Department of Nuclear Engineering, University of California); Dr KHESWA, Bongani (Department of Physics, University of Oslo); Dr BLEUEL, Darren (Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory); Dr BURKE, Jason (Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory); Dr RESSLER, Jo (Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory); Dr KOGLIN, John (Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory); Dr BERNSTEIN, Lee (Nuclear Science Division, Lawrence Berkeley National Laboratory); Dr CRABTREE, Lily (Department of Nuclear Engineering, University of Tennessee); Dr PETRI, Maria (Nuclear Science Division, Lawrence Berkeley National Laboratory); Dr SCIELZO, Nick (Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory); Dr ORCE, Nico (University of the Western Cape); Dr PAPKA, Paul (Stellenbosch University); Dr HATARIK, Robert (Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory); Dr PASCHALIS, Stefanos (Nuclear Science Division, Lawrence Berkeley National Laboratory); Dr SIEM, Sunniva (Department of Physics, University of Oslo); Dr REED, Tim (Department of Nuclear Engineering, University of Tennessee)

Presenter: Mr SOWAZI, Khanyisa (iThemba LABS)

Contribution ID: 12

Type: **not specified**

Coupling of single neutron and proton configurations to collective core excitations in ^{162}Yb .

The excited states of ^{162}Yb have been studied at iThemba Laboratory for Accelerator Based Sciences (iThemba LABS), using the $^{150}\text{Sm} (^{16}\text{O}, 4n) ^{162}\text{Yb}$ fusion-evaporation reaction. The beam of 83 MeV ^{16}O was provided by the Separated-Sector Cyclotron (SSC) and used to bombard a 3 mg/cm^2 ^{150}Sm target. The gamma rays emitted from the reaction products were detected using the AFRODITE gamma-ray spectrometer, comprised of 8 Compton-suppressed clover detectors. Attempts have been made in identifying the low-lying positive parity bands in ^{162}Yb , particularly the beta and gamma vibrational bands, which are traditionally associated with the first excited $K^\pi = 0^+$ and $K^\pi = 2^+$ respectively. Many levels have been found. In particular the first excited 0^+ band and the even and odd sequences of the gamma band have been firmly established. The 0^+ band and the even spin members of the gamma band are observed to exhibit a Landau-Zener crossing. This crossing demonstrates that the signature splitting in gamma bands is mainly caused by band mixing. The data will be discussed in terms of the Triaxial Projected Shell Model and also with the predictions of the 5-Dimensional Collective Model (5-DCM).

Primary author: Mr MDLETSHE, Linda (University of Zululand)

Co-authors: Prof. SHARPEY-SCHAFER, J. F (University of the Western Cape); Mr SITHOLE, M. A (University of the Western Cape); Ms KHUMALO, N. A (University of the Western Cape); Dr BARK, R. A (iThemba LABS); Ms JONGILE, S (University of Zululand); Dr MAJOLA, S. N. T (University of Zululand); Dr NTSANGASE, S. S (University of Zululand); Dr DINOKO, T. S (iThemba LABS)

Presenter: Mr MDLETSHE, Linda (University of Zululand)

Contribution ID: 13

Type: **not specified**

Search for Charality in ^{193}Tl

Research conducted at iThemba LABS showed that chiral symmetry can develop in the thallium isotopes in the 190 mass region. In order to increase the knowledge about chirality in this mass region, a γ -spectroscopy study of ^{193}Tl was performed at iThemba LABS. The previous level scheme of ^{193}Tl was modified and extended. Spin and parity were assigned to most of the levels. Three negative parity bands showing similar properties were identified. These bands were associated with the same configuration which is suitable for chiral symmetry. The observed near-degeneracy is good and indicates the presence of chiral symmetry. Furthermore, two bands that could form a chiral pair were observed at higher spins. The results from theoretical calculations using the Cranked Nilsson-Strutinsky (CNS) codes and the multi-particle-plus-triaxial rotor (MPR) model of Carlsson and Ragnarsson are in agreement with the proposed observation of chiral symmetry. Possible multiplet of chiral systems will be discussed.

Primary author: Dr NDAYISHIMYE, Joram (iThemba LABS)

Co-author: Dr LAWRIE, Elena (iThemba LABS)

Presenter: Dr NDAYISHIMYE, Joram (iThemba LABS)

Contribution ID: 14

Type: **not specified**

Tape station for beta decay measurement

The tape station system is used for collecting and counting of “mostly” short-lived radioactive nuclei and these kinds of systems are used at an on-line facility. Such devices allow measuring radioactive particle even if they cannot be seen by other instruments. A flexible tape transport system may be used to catch recoiling nuclei after they have been mass identified by high-resolution mass spectrometer.

Depending on the half-life of the element of interest, where the detection point can be same as the implantation point or measuring at the different position, a tape has to be moved to a low background area where it will be observed by and detectors. After counting for a while, the tape is moved again to make sure that activity from the complete decay chain to stability is not observed.

Primary author: Mr MAKHATHINI, Lucky (iThemba LABS)

Co-author: Dr BARK, Rob Bark (iThemba LABS)

Presenter: Mr MAKHATHINI, Lucky (iThemba LABS)

Contribution ID: 15

Type: **not specified**

Coupling of single proton configurations to collective core excitations in ^{162}Yb : the nucleus ^{161}Tm

Over the past two decades there has been controversy as to whether deformed nuclei are subject to quadrupole vibrations (γ and β), particularly the β vibrations. Pertaining the $\gamma(K=2^+)$ vibrations, experimental evidence has been more or less consistent, confirming they indeed exist. On the other hand the situation remains elusive for the β vibrations which are characterized by the first low lying 0^+ excited state. The current study seeks to get more insight on the microscopic nature of the aforementioned by studying the nucleus ^{161}Tm , which was populated using the $^{152}\text{Sm}(^{14}\text{N}, n)^{161}\text{Tm}$ reaction with the aid of the AFRODITE array at iThemba LABS. A level scheme was built for ^{161}Tm by examining multiple gates using coincidence spectra. Transitions were confirmed with DCO (Direct Correlations for Oriented states) and/or polarization anisotropy measurements where applicable. Alignments and band crossings have been used to meaningfully describe the quantum behavior of the collective structures observed in this work. In addition systematic comparisons have also been used to further understand the structural behavior of band structures observed in the level scheme. Furthermore experimental $B(M1)/B(E2)$ values for bands involving the $[505]11/2^-$ orbital and other observed strong coupled bands were obtained to confirm quasi-particle configurations.

Primary author: Ms JONGILE, SANDILE (UNIVERSITY OF ZULULAND)

Co-authors: Mr SITHOLE, ABEL MAKUHANE (UNIVERSITY OF THE WESTERN CAPE); Prof. SHARPEY-SCHAFFER, JOHN (UNIVERSITY OF THE WESTERN CAPE); Mr MDLETSHI, LINDA (UNIVERSITY OF ZULULAND); Dr NTSHANGASE, SIFISO SENZO (UNIVERSITY OF ZULULAND); Dr MAJOLA, SIYABONGA (UNIVERSITY OF ZULULAND); Dr DINOKO, TSHEPO (ITHEMBA LABS)

Presenter: Ms JONGILE, SANDILE (UNIVERSITY OF ZULULAND)

Contribution ID: 16

Type: **not specified**

LOW SPIN STRUCTURE STUDIES OF EXOTIC STATES AND NUCLEI: NEW CAPABILITIES AT ITHEMBA LABS

Excited 0^+ states are the least understood of any low-energy degree of freedom in nuclei. $E0$ transition strengths are a measure of the off-diagonal matrix elements of the mean-square charge radius operator. Mixing of configurations with different mean-square charge radii produces $E0$ transition strength. The study of low-lying 0^+ states has been a focus of attention in the last two decades because of their implications regarding shape coexistence in nuclei.

New capabilities are underway to be added to the arsenal of cutting-edge spectroscopy techniques at iThemba LABS. Through the application of electron and pair spectroscopy, and with the addition of magnetic spectrometers new avenues of nuclear structure will be opened and explored.

In addition, an array of LaBr₃(Ce) detectors (2"x2") for fast-timing measurements have been commissioned. With excellent timing resolution (<350ps) and good energy resolution these will be used to study the decays of the photon branch of these states. Measurements of lifetimes of these states, and their associated feeding/decaying states are also extremely important in understanding the associated wave functions and orbital assignments of states. With the combination of an array of fast-timing detectors, this enhances the motivation for these nuclear structure studies over a region of nuclear structure.

A review of these capabilities will be presented and commissioning experiments presented.

Primary author: Dr JONES, Pete (iThemba LABS)

Co-authors: Mr AVAA, Abraham (iThemba/Wits); Mr ZIKHALI, Bonginkosi (iThemba/UWC); Mr MSEBI, Lumkile (iThemba/UWC); Mr INGERBERG, Vetle (iThemba LABS / University of Oslo)

Presenter: Dr JONES, Pete (iThemba LABS)

Contribution ID: 17

Type: **not specified**

The spectroscopy of ^{162}Hf at low and high spins

The search for triaxial strongly deformed (TSD) structures in the rare earth region of the nuclear chart has been of interest recently. These structures have been predicted to occur for over 35 years [1]. The main aim of this project is to comprehensively study the nuclear spectroscopy of ^{162}Hf at low and high spins and to search for TSD structures. The research is also part of an investigation into the systematic behaviour of the N=90 nuclei. Our interest is in the systematics of the positive parity excited bands in N=90 nuclei and the behaviour of the negative parity bands as the proton number Z is increased leading to a reduction in the deformation of the ground state structure. A number of high spin structures with high dynamic moments of inertia have been observed in the heavy Hafnium isotopes [2-7]. Cranked Nilsson Strutinsky calculations predict that such bands are most likely associated with TSD structures originating from a positive- γ energy minimum which dominates at ultrahigh spins. The data was collected using the world class multi-detector gamma-ray spectrometers namely the AFRODITE (South Africa) and JUROGAM at Jyvaskyla (Finland). The high spin states were populated in the $^{110}\text{Pd}(^{56}\text{Fe}, 4n)$ reaction at a beam energy of 270 MeV at Jyvaskyla and the medium and low spin states at iThemba LABS using the $^{144}\text{Sm}(^{22}\text{Ne}, 4n)$ reaction at a beam energy of 110 MeV. The collected data was sorted using MTSort and was analyzed using RADWARE [8]. Results from both data sets will be presented.

- [1] G. Andersson et al., *Nucl. Phys. A* 268, 205 (1976)
- [2] H. Amro et al., *Phys. Lett. B* 506, 39 (2001)
- [3] A. Neußer-Neffgen et al., *Phys. Rev. C* 73, 034309 (2006)
- [4] Y. C. Zhang et al., *Phys. Rev. C* 76, 064321 (2007)
- [5] D. J. Hartley et al., *Phys. Lett. B* 608, 31 (2005)
- [6] D. T. Scholes et al., *Phys. Rev. C* 70, 054314 (2004)
- [7] R. B. Yadav et al., *Phys. Rev. C* 78, 044316 (2008)
- [8] D.C Radford. *ESCL8R and LEVIT8R: Nuclear Instruments and Methods in Physics Research A*, 361: 297-305, January (1995)

Primary author: Ms KHUMALO, Nontobeko (University of the Western Cape)

Co-authors: Mr ZIKHALI, Bonginkosi (University of Zululand); LAWRIE, Elena (iThemba LABS); Prof. SHARPEY-SCHAFFER, John F (University of Western Cape); Dr NDAYISHIMYE, Joram (iThemba LABS); Mr MALATJI, Kgashane (University of the Western Cape); Dr LAWRIE, Kobus (iThemba LABS); Mr MDLETSHE, Linda (University of Zululand); Dr SHIRINDA, Obed (iThemba LABS); Dr JONES, Pete (iThemba LABS); Dr BARK, Robert (iThemba LABS); Dr NCHODU, Rudolph (iThemba LABS); Ms JONGILE, SANDILE (UNIVERSITY OF ZULULAND); Dr MAJOLA, SIYABONGA (UCT/ITHEMBA LABS); Dr NTSHANGASE, Sifiso Senzo (University of Zululand); Mr NONCOLELA, Sive (UWC); Dr BUCHER, Thifhelimbilu Daphney (iThemba LABS); Dr DINOKO, Tshepo (University of the Western Cape/ iThemba LABS); Mr SITHOLE, makuhane (University of Venda); Dr WIEDEKING, mathis (itl)

Presenter: Ms KHUMALO, Nontobeko (University of the Western Cape)

Contribution ID: 18

Type: **not specified**

A review of nuclear cluster models

The nuclear cluster models has extensively been used to understand some of the structural properties of nuclear matter. A number of theoretical models have been developed since the discovery that alpha decay could be conceptually understood as the expulsion of a correlated subset of four nucleons from the parent nucleus. This presentation will give a brief overview of the most popular nuclear cluster model descriptions in nuclear physics, and will highlight some of the strengths and weaknesses of each of the models.

Primary author: Dr WYNGAARDT, Shaun (Stellenbosch University)

Co-authors: Mr KIMENE KAYA, Christel (Stellenbosch University); Dr IBRAHIM, taofiq (Stellenbosch University)

Presenter: Dr WYNGAARDT, Shaun (Stellenbosch University)

Contribution ID: 19

Type: **not specified**

Study of the low-lying states in superscript²⁶ Mg nuclei.

The study of ²⁶Mg level structure is of particular importance in nuclear structure physics because of its impact in the slow neutron capture process (s-process) for the nucleosynthesis of heavy elements up to lead. In fact, the ²⁶Mg is the compound nucleus for the s-process neutron source ²²Ne (α, n)²²Mg that is one of the dominant reaction in the s-process. Since it is an endothermic reaction it competes directly with the ²²Ne (α, γ)²⁶Mg radiative capture. Understanding the rate of both of these reactions is crucial for linking the observational evidence of s-process abundance with the internal structure of the stars. The uncertainties in the energy of the ²⁶Mg states and the inconclusive spin-parity assessments still lead in large error bars in these reaction rates.

In this work, the low-energy states in ²⁶Mg were populated using the inelastic scattering of alpha particles with beam energy of 120 MeV. The ²⁶Mg (α, α')
²⁶Mg measurements can be useful to improve the uncertainties in these rates by studying and extracting the characteristics of the excited states of ²⁶Mg. The experiment was performed at iThemba LABS research facility using a new designed experimental set-up: the K600 magnetic spectrometer coupled to the BaGeL array (Ball of Germanium and LaBr detectors). This combination was used to perform the coincidence measurements between charged particles and gamma rays. The excitation and the subsequent gamma decay of the ²⁶Mg states of interest will be investigated. Preliminary results on the analysis will be presented.

Primary authors: Dr MARIN-LAMBARRI, Daniel (iThemba LABS); Prof. SIDERAS-HADDAD, Elias (Witswatersrand); Ms JIVAN, Harshna (University of the Witwatersrand); Prof. CARTER, John (School of Physics, Wits University); Dr PELLEGGRI, Luna (iThemba LABS & WITS); Dr PAPKA, Paul (Stellenbosch University); Ms MOLEMA, Pheladi (witwatersrand); Dr NEVELING, Retief (iThemba LABS); Dr SMIT, Ricky (iThemba LABS); Dr PESUDO, Vicente (iThemba LABS); Mrs KHESWA, ntombi-zonke (iThemba LABS)

Presenter: Ms MOLEMA, Pheladi (witwatersrand)

Contribution ID: 20

Type: **not specified**

Measurements of natural radioactivity in sands using an array of lanthanum -bromide scintillator detectors

LaBr₃:Ce detectors have been shown to be 1.2–1.65 times more efficient than NaI:Tl detectors above 350 keV, for 3.8 cm×3.8 cm (1.5 in.×1.5 in.) detectors and have an energy resolution of 2.5–3% at the 662 keV gamma-line of ¹³⁷Cs, compared to 6–7% for NaI:Tl detectors[1]. The detector crystal has other advantages such as a high scintillation light output with a fast decay time[2]. An array of 8 2in x 2in LaBr₃(Ce) scintillators with an XIA PIXIE-16 Digital Signal Processing system data acquisition system will be used to measure sands placed in the centre (17.5 cm from each detector) of the array (with all detectors lying in the horizontal plane) for 12 hours. The gamma-gamma coincidence method has the advantage of virtually eliminating all background peaks that do not exist in coincidence with other peaks, significantly improving detection limits of useful radionuclides[3][4]. By employing a gamma-gamma coincidence condition, the background from the radioisotopes in the LaBr₃:Ce scintillator is eliminated, providing a means for improving detection limits[5]. The absolute gamma-ray energy detection efficiency of each detector will be determined and compared. Data from each detector will be analyzed. The activity concentration of ²³⁸U, ²³²Th and ⁴⁰K in the sands will be determined and compared to certified values. Radiation hazard indices will be calculated for the soil samples and compared to certified values.

Reference

- [1] K. Ciupek, S. Jednoróg, M. Fajak, and K. Szewczak, “Evaluation of efficiency for in situ gamma spectrometer based upon cerium-doped lanthanum bromide detector dedicated for environmental radiation monitoring,” *J. Radioanal. Nucl. Chem.*, vol. 299, no. 3, pp. 1345–1350, 2014.
- [2] A. Favalli, H. C. Mehner, and F. Simonelli, “Wide energy range efficiency calibration for a lanthanum bromide scintillation detector,” *Radiat. Meas.*, vol. 43, no. 2–6, pp. 506–509, 2008.
- [3] M. Yoho and S. Landsberger, “Determination of Selenium in coal fly ash via γ - γ coincidence neutron activation analysis,” *J. Radioanal. Nucl. Chem.*, vol. 307, no. 1, pp. 733–737, Jan. 2016.
- [4] S. Horne and S. Landsberger, “Selenium and mercury determination in biological samples using gamma-gamma coincidence and Compton suppression,” *J. Radioanal. Nucl. Chem.*, vol. 291, no. 1, pp. 49–53, Jan. 2012.
- [5] A. Drescher et al., “Gamma-gamma coincidence performance of LaBr₃:Ce scintillation detectors vs HPGe detectors in high count-rate scenarios,” *Appl. Radiat. Isot.*, vol. 122, no. January, pp. 116–120, 2017.

Primary author: Ms BASHIR, Munirat (Stellenbosch University/iThemba)

Co-authors: Dr PETE, Jones (iThemba LABS); Prof. NEWMAN, R.T (Stellenbosch University)

Presenter: Ms BASHIR, Munirat (Stellenbosch University/iThemba)

Contribution ID: 22

Type: **not specified**

Spectroscopy of quadrupole excitations at low spin in even-even $N \sim 90$ nuclei

Vibrational levels are well known in atomic nuclei but despite many decades of research, some of their properties still remain elusive. In particular, low-lying rotational bands based on the first excited 0^+ state, which are traditionally understood as β vibrational bands nevertheless show properties at odds with this interpretation, more especially in the transitional rare earth region with $N \sim 90$ [Gar01]. An alternative is that they can be better described as a “second vacuum”, or coexisting minimum in the pairing degree of freedom [Sch11a]. In order to produce a complete and definitive microscopic picture of the so-called β and γ bands, an extensive systematic review is performed for nuclides in the 160 mass region, between $N = 88$ and 92 and Sm to Yb. The data are explained using a five dimensional collective Hamiltonian for quadrupole rotational and vibrational degrees of freedom [Li09, Nik09]. A good qualitative agreement is obtained between measured energies and electromagnetic transition rates across the entire $A \sim 160$ mass region. The implication of these findings on the interpretation the first excited 0^+ states is there from discussed.

[Gar01] P. E. Garrett, J. Phys. G27, R1 (2001)

[Li09] Z.P. Li et al., Phys. Rev. C 79, 054301 (2009)

[Nik09] T. Nikšić., Phys. Rev. C 79, 034303 (2009)

[Sch11a] J.F. Sharpey-Schafer et al. Eur. Phys. J. A47 (2011) 6

Primary author: Mr MAJOLA, SIYABONGA (ITHEMBA LABS/UNIZULU)

Co-authors: Mr ZIMBA, George (University of Johannesburg); Prof. SHARPEY-SCHAFER, John F (University of Western Cape); Mr STANKIEWICZ, Maciej (UCT); Dr BARK, Robert (iThemba LABS); Ms BVUMBI, SUZAN PHUMUDZO (UNIVERSITY OF JOHANNESBURG); Mr ZHI, Shi (Peiking University); Dr ZHANG, Shuangquan (School of Physics, Peking University, Beijing); Dr NTSHANGASE, Sifiso Senzo (University of Zululand); Mr DINOKO, Tshepo (University of the Western Cape/ iThemba LABS); Prof. LI, Zhipan (South West Univesity)

Presenter: Mr MAJOLA, SIYABONGA (ITHEMBA LABS/UNIZULU)

Contribution ID: 23

Type: **not specified**

Multiple many-particle chiral systems described within the particle-rotor model

A nuclear chiral system is formed when the total angular momentum of the nucleus is aplanar, i.e. when it has significant projections along all three nuclear axes [1]. Most important for the identification of chiral bands is to establish a pair of $\Delta I = 1$ bands that are near-degenerate in energy, but also in B(M1) and B(E2) transition probabilities [1]. Up to date, chiral candidates showing two- or multi-quasiparticle partner bands have been observed in several nuclei in $A \sim 80, 100, 130$ and 190 mass regions. It was suggested that multiple chiral partner bands (MXD) with the same two-quasiparticle nucleon configuration may exist in a single nucleus [1-3]. The first multiplet of chiral bands built on the same multi-quasiparticle configuration was observed in ^{103}Rh [4]. Such a scenario was also considered recently for ^{194}Tl [5].

In this work the multi-particle-plus-triaxial-rotor (MPR) [6] model calculations were performed for multiple chiral systems associated with the same multi-quasiparticle configurations in the 100, 130 and 190 mass regions described with a realistically large configuration space for the odd proton(s) and the odd neutron(s). The main objective was to study the properties of the excited chiral systems associated with the same nucleon configuration. In particular we were interested in whether the excited chiral systems might show better near-degeneracy than the yrast one.

Multiple chiral systems associated with many-particle nucleon configurations were found, but they may not necessarily form pairs of near-degenerate bands in an obvious way. Our calculations showed that in order to search for the best chiral symmetry, one needs to study not only the two lowest-energy bands, but also as many excited bands as possible. It is quite possible that the excited bands will couple into pairs with more similar geometry of the intrinsic angular momenta as a function of spin, and show closer intrinsic structure, than the two lowest-energy bands. It is also concluded that to couple chiral bands into pairs with a similar nature one needs to consider the projections of the angular momenta along the nuclear axes. Part of the results from this work has been published in Ref. [7]. The results from these calculations will be presented and discussed.

This work is based upon research supported by the National Research Foundation, South Africa, with grant numbers 75357, 76632, 91446, 93531, and 103478. We thank B.G. Carlsson and I. Ragnarsson for making available the multi-particle-plus-triaxial-rotor model codes and for numerous fruitful discussions.

[1] S. Frauendorf, J. Meng, Nucl. Phys. A 617, 131 (1997).

[2] Q.B. Chen et al., Phys. Rev. C 82, 067302 (2010).

[3] H. Zhang, Q. Chen, Chin. Phys. C 40, 024102 (2016).

[4] I. Kuti et al., Phys. Rev. Lett. 113, 032501 (2014).

[5] P.L. Masiteng et al., Eur. Phys. J. A 50, 119 (2014); P.L. Masiteng et al, Eur. Phys. J. A 52, 28 (2016).

[6] B.G. Carlsson, I. Ragnarsson, Phys. Rev. C 74, 044310(2006); B.G. Carlsson, Models for rotating nuclei - cranking and rotor plus particles coupling, PhD Thesis (Lund University, 2007) ISBN 978-91-628-7355-4; B.G. Carlsson, Int. J. Mod. Phys. E 16, 634 (2007).

[7] O. Shirinda and E.A. Lawrie, Acta Phys. Pol. B46, (2015) 683; O. Shirinda and E.A. Lawrie, Eur. Phys. J. A52, 344 (2016).

Primary author: Dr SHIRINDA, Obed (iThemba LABS/Stellenbosch University)

Co-author: Dr LAWRIE, Elena (iThemba LABS)

Presenter: Dr SHIRINDA, Obed (iThemba LABS/Stellenbosch University)

Contribution ID: 24

Type: **not specified**

Nuclear structure studies relevant to double beta decay of ^{136}Xe

In addition to establishing the Majorana nature of neutrinos, obtaining the absolute neutrino mass scale is now the focus of several large-scale neutrinoless double beta decay experiments.

The current challenge in determining the neutrino mass accurately depends on calculation of nuclear matrix elements (NME's) in the select nuclei where these decays can take place. It is well known that the dominating uncertainties in the calculated NME values arise from the model dependence of these calculations.

In this talk we will present some recent experimental results using high resolution spectroscopy from $^{136,138}\text{Ba}(p,t)$ and $^{138}\text{Ba}(d,a)$ reactions that will be useful for future NME calculations for the double beta decay of $^{136}\text{Xe} \rightarrow ^{136}\text{Ba}$

Primary author: Ms REBEIRO, BERNADETTE (UNIVERSITY OF THE WESTERN CAPE, SOUTH AFRICA)

Co-authors: Ms DIAZ VARELA, Alejandra (University of Guelph); Ms RADICH, Allison (University of Guelph); Mr JIGMEDDORJ, Badamsambuu (University of Guelph); Ms BURBADGE, Christina (University of Guelph); Mr RAND, Evan (University of Guelph); Mr BALL, Gordon (TRIUMF); Mr WIRTH, Hans (Ludwig-Maximilians-Universitat Munchen); Mr ONDZE, Jespere (UNIVERSITY OF THE WESTERN CAPE); Mr LEACH, Kyle (Colorado School of Mine); Mr GARRETT, Paul (University of Guelph); Mr ADSLEY, Philip (Themba Laboratory for Accelerator Based Sciences, University of Stellenbosch, Nucleaire d'Orsay); Mr HERTENBERGER, Ralph (Ludwig-Maximilians-Universitat Munchen); Prof. LINDSAY, Robbie (UWC); TRIAMBAK, Smarajit (University of Western Cape); Mr FAESTERMANN, Thomas (Technische Universitat Munchen); Mr BILDSTEIN, Vinzenz (University of Guelph); Ms MABIKA, Zandile (University of the Western Cape)

Presenter: Ms REBEIRO, BERNADETTE (UNIVERSITY OF THE WESTERN CAPE, SOUTH AFRICA)

Contribution ID: 25

Type: **not specified**

Radiation Safety Calculations for iThemba LABS ACE Isotopes facility design - Faraday Cups

The design of iThemba LABS Accelerator Centre for Exotic (ACE) Isotopes facility was proposed to have four beam-lines feeding the beam to two production vaults, with each vault having two production stations. The plan is to simultaneously operate one production station in both production vaults, meaning that only two opposite beam-lines will be delivering the beam at a time, with each of the two beam-lines feeding 70 MeV energy proton-beam of up to 350 μA current.

Two Faraday cups will be simultaneously used to stop 50 μA (standard beam current) proton beams during the optimization of beam-lines parameters.

The whole ACE operating system will be accompanied by an interlock system that will switch off the parameters of the cyclotron, e.g. the RF, and insert a Faraday cup in the axial injection line in case the beam is accidentally stopped in the beam-line. Moreover, less than 5% beam loss is expected to occur in the cyclotron. Beam-pipes of 150 mm diameter were proposed, therefore, the beam losses in beam-lines are not expected to be significant. Radiation resulting from beam loss in the cyclotron (35 μA at most) will, to a large extent, be shielded by the magnet of the cyclotron. The magnet shielding of the radiation caused by beam loss inside the cyclotron is better in the vertical direction. Therefore, radiation that will make it through the cyclotron vault floor to the basement is not expected to be significant.

Consequently, for the shielding of the cyclotron, the main source of radiation to consider was the beams on Faraday cups; Hence, this study was looking at the optimal shielding design especially when the proton beams hit the Faraday cups during optimization of the cyclotron beam-line parameters. This study was conducted using the FLUKA Monte-Carlo transport code.

Primary author: Mr MANTENGU, Nkanyiso Robert (iThemba LABS)

Co-author: Mr BEUKES, Philip (iThemba LABS)

Presenter: Mr MANTENGU, Nkanyiso Robert (iThemba LABS)

Contribution ID: 26

Type: **not specified**

New developments in the nuclear binary cluster-Core in the Super-Heavy Nuclear region

The atomic nucleus is a complex many-body interacting system, which exhibits a underlying correlated set of nucleon states. The clustering model is one of the most reliable models that predicts the strongly correlated subsystem of nucleons closed to the threshold decay of nuclei. The binary-cluster model describes the structure and decay properties of super-heavy nuclei.

The phenomenological Cubic Woods-Saxon potential, developed by Buck, Merchant and Perez, has successfully predicted a number of experimental observables associated with clustering phenomenon. The recently developed microscopic double folded M3Y potential results in the inverted spectra for the positive parity excited cluster states, but successfully predicts the decay half-life for alpha-Pb system. These shortcomings of the M3Y based microscopic binary cluster model lead to the newly developed hybrid code-cluster potential, obtained by fitting the Phenomenological Woods-Saxon Cube and the M3Y double folding at the surface region where the two potential coalesce.

The project presented will give an overview of the nuclear cluster models available. Furthermore the recently developed self-consistent relativistic mean-field cluster-core description will be presented.

Primary author: Mr KIMENE KAYA, Christel (Stellenbosch University)

Co-author: Prof. WYNGAARDT, Shaun (Stellenbosch)

Presenter: Mr KIMENE KAYA, Christel (Stellenbosch University)

Contribution ID: 27

Type: **not specified**

Coulomb-excitation programme at UWC

The Coulomb-excitation process with the combination of highly-efficient γ and particle detector systems present a great tool to study quadrupole collectivity in nuclei and probing nuclear-structure properties. This process employs the well-known Coulomb interaction and selectively excites collective nuclear states which decay back to the ground state through γ -ray transitions. Coulomb-excitation measurements have been carried out by the UWC Coulex group, where various reorientation-effect Coulomb-excitation measurements (RECE) have recently been carried out f

or a systematic study throughout nuclei in the sd shell. This work reports on our new measurements on $Q_s(2_1^+)$ values at iThemba LABS and TRIUMF on ($^{12}C, ^{20}Ne, ^{32}S, ^{36}Ar$ and ^{40}Ar), which will be presented during this conference. In particular, a solution is proposed for the striking zig-zag pattern of $Q_s(2_1^+)$ values observed at the end of the sd shell.

Primary author: Ms MASANGO, Senamile (University of the Western Cape)

Presenter: Ms MASANGO, Senamile (University of the Western Cape)

Contribution ID: 28

Type: **not specified**

Spectroscopy of Exotic Lambda-hypernuclei within Covariant Density Functional Theory

Covariant density functional theory can well describe the bulk and single-particle properties of nuclei in a self-consistent way. Typical neutron-rich Lambda-hypernuclei ^{12}Be and ^{16}C are investigated within covariant density functional theory. The bulk properties of mean field, meson-nucleon tensor coupling, spin-orbit term, behaviors of neutrons are focused.

Primary author: LU, Hongfeng (College of Science, China Agricultural University)

Presenter: LU, Hongfeng (College of Science, China Agricultural University)

Contribution ID: 29

Type: **not specified**

Multiple Chiral Doublet Bands in ^{126}Cs

Following the previously observed chiral doublet bands in ^{126}Cs , a new pair of chiral doublet bands has been proposed. Both two pairs are based on the identical configuration $\pi h_{11/2} \otimes \nu h_{11/2}$ and may be another candidate of novel type of multiple chiral doublets (“yrast” and “excited” $M\chi D$) first observed in ^{103}Rh . Theoretical calculations within covariant density functional theory and particle rotor model are also performed.

Primary authors: Dr LI, Jian (College of physics, jilin university); Mr LIU, Jiaqiang (College of Physics, Jilin University, Changchun 130012, China); Prof. LU, Jingbin (College of Physics, Jilin University, Changchun 130012, China); Mr LIU, Yonghao (College of Physics, Jilin University, Changchun 130012, China)

Presenter: Dr LI, Jian (College of physics, jilin university)

Contribution ID: 30

Type: **not specified**

Multipole Response of Hypernuclei within Skyrme Energy Density Functional

We have extended the Hartree-Fock and Random Phase Approximation method to calculate the multipole giant resonances of hypernuclei within the Skyrme energy density functional self-consistently. This means that the same Skyrme interactions, including the nucleon-nucleon, nucleon-hyperon as well as hyperon-hyperon interaction, are adopted in the calculation of the properties of ground states and collective states in hypernuclei. In the calculations we use SKM* Skyrme interaction for the nucleon-nucleon interaction and Yamamoto nucleon-hyperon interaction as well as the Lanskoy $S\Lambda\Lambda 1$ parameter set for hyperon-hyperon interaction. We have systematically investigated the multipole response of double- Λ hypernuclei $^{18}\text{O}\Lambda\Lambda$, $^{42}\text{Ca}\Lambda\Lambda$, $^{92}\text{Zr}\Lambda\Lambda$, $^{122}\text{Sn}\Lambda\Lambda$ and $^{210}\text{Pb}\Lambda\Lambda$. The effect of hyperon interaction on response function, the transition density, the low-lying states are analyzed in more details, the role of hyperon interaction depends on the type of the collective excitations, in some case it gives attractive effect, in another case, it is repulsive.

Primary author: Prof. CAO, Ligang (School of Mathematics and Physics, North China Electric Power University, Beijing 102206, China)

Co-author: Mrs LV, Hong (School of Mathematics and Physics, North China Electric Power University, Beijing 102206, China)

Presenter: Prof. CAO, Ligang (School of Mathematics and Physics, North China Electric Power University, Beijing 102206, China)

Contribution ID: 31

Type: **not specified**

Exploration of direct neutron capture with covariant density functional theory inputs

Predictions of direct neutron capture are of vital importance for simulations of nucleosynthesis in supernovae, merging neutron stars, and other astrophysical environments. We calculated direct capture cross sections using nuclear structure information obtained from a covariant density functional theory as input for the FRESKO coupled reaction channels code. We investigated the impact of pairing, spectroscopic factors, and optical potentials on our results to determine a robust method to calculate cross sections of direct neutron capture on exotic nuclei. Our predictions agree reasonably well with experimental cross section data for the closed shell nuclei ^{16}O and ^{48}Ca , and for the exotic nucleus ^{36}S . We then used this approach to calculate the direct neutron capture cross sections on Sn isotopes which are of interest for the astrophysical r-process.

Primary author: Dr SHISHENG, Zhang (Beihang University)

Presenter: Dr SHISHENG, Zhang (Beihang University)

Contribution ID: 35

Type: **not specified**

Improving of the injection line transmission through the SPC2 cyclotron at iThemba LABS

iThemba LABS has recently engaged in a journey of producing metal ion beams to use for nuclear physics experiments. In order to successfully deliver these beams, it is important that the transmission of the injection beamline through the injector cyclotron (SPC2) is of high quality. In order to achieve this, efforts are being made to improve the transmission. One of the concepts being investigated is how adding a second buncher, operating in second harmonic will improve the transmission. This is done using different simulation programs. In this talk preliminary results of the simulations will be presented.

Primary author: Mr NEMULODI, FHUMULANI (ITHEMBA LABS)

Co-authors: Mr FOURIE, Dirk (iThemba LABS); Mr WILLIAM, Duckitt (iThemba LABS); Mr DE VILLIERS, Garrett (iThemba LABS); Dr YANG, Jingjuan (CIAE); Mr MIRA, Joele Paulus (iThemba LABS/University of Stellenbosch); Mr VAN NIEKERK, Johan (iThemba LABS); Dr CONRADIE, Lowry (iThemba LABS); Dr THOMAE, Rainer (iThemba LABS); Dr ZHANG, Tianjue (CIAE)

Presenter: Mr NEMULODI, FHUMULANI (ITHEMBA LABS)

Contribution ID: 36

Type: **not specified**

Measurement of beam characteristics of Low-Energy Accelerators

China JinPing underground Laboratory (CJPL) was established inside the tunnels piercing Jinping Mountain in Sichuan Province, China, which can provide an ideal environment for low background experiment. A new 400 kV accelerator, with high current based on an ECR source, will be placed at this underground laboratory for nuclear astrophysics experiment (JUNA). Due to the new accelerator still being under construction, the resonance reactions, like $^{27}\text{Al}(p, \gamma)^{28}\text{Si}$ and $^{24}\text{Mg}(p, \gamma)^{25}\text{Al}$, and non-resonance $^{12}\text{C}(p, \gamma)^{13}\text{N}$ was studied at 320 kV high-voltage platform at IMP to estimate the proton beam characteristics, like absolute energy, energy spread, and long-term energy stability. The results of experiment and current state of new 400 kV accelerator will be given.

Primary authors: Dr LI, KUOANG (IMP,CAS,CHINA); Dr WANG, SHUO (SHANDONG UNIVERSITY)

Presenter: Dr WANG, SHUO (SHANDONG UNIVERSITY)

Contribution ID: 38

Type: **not specified**

Possible umbrella-like antimagnetic rotation (UAMR) mode in Pd isotopes

The tilted axis cranking model based on covariant density functional theory is used to study the $\pi g_{9/2}^{-4} \nu h_{11/2}$ bands in odd-A $^{101,103}\text{Pd}$ and the $\pi g_{9/2}^{-4} \nu h_{11/2}^2$ bands in even-even $^{102,104}\text{Pd}$. The experimental energy spectra and $B(E2)$ values are reproduced well in the self-consistent and microscopic calculations. By investigating microscopically the composition and orientation of angular momentum, an umbrella-like antimagnetic rotation (UAMR) mode resulting from the coupling of four $g_{9/2}$ proton holes to one or two aligned $h_{11/2}^2$ neutron particles is clearly illustrated for the first time.

Primary author: Dr BIN, Qi (Shandong University, Weihai)

Presenter: Dr BIN, Qi (Shandong University, Weihai)

Contribution ID: 39

Type: **not specified**

Chiral bands in ^{82}Br

High spin states of ^{82}Br were studied by using the fusion-evaporation reaction $^{82}\text{Se} (^4\text{He}, 1\text{p}4\text{n})$ at beam energies of 65 and 68 MeV. 26 new transitions and 13 new levels were added into the previous level scheme, including a positive-parity band. With the earlier configuration assignment $\pi 9/2 \otimes \nu 9/2$ to band 1 and the similar experimental features to the chiral doublet bands observed in ^{78}Br and ^{80}Br , the positive-parity doublet bands 1 and 2 in ^{82}Br are therefore suggested as chiral doublet bands with the $\pi 9/2 \otimes \pi 9/2$ configuration. The interpretation was supported by the relativistic mean field theory and triaxial particle rotor model calculations.

Primary author: Mr GUO, Ruiju (Shandong University at Weihai)