## First Pan-African Astro-Particle and Collider Physics Workshop



Contribution ID: 1

Type: not specified

## Laser-assisted processes beyond the standard model

Monday, 21 March 2022 15:10 (15 minutes)

In this work, we have theoretically studied the neutral Higgs pair production in Two Higgs Doublet Model (THDM) in the presence of a circularly polarized laser field. The laser-assisted differential partial cross section is derived in the centre of mass frame at the leading order including Z diagram. The total cross section is computed numerically by integrating the differential cross section over the solid angle d $\Omega$ . Two benchmark points are discussed for the THDM parameters. In the first step, we have analyzed the total cross section of e+e-  $\rightarrow$  h0A0 by considering H 0 as the standard model-like Higgs boson. Then, the process e+e-  $\rightarrow$  H0A0 is studied by taking h0 as the Higgs boson of the standard model. For both benchmark points, the laser-assisted total cross section of the studied processes depends on the produced neutral Higgs masses, the centre of mass energy and the laser field parameters. In addition, the maximum cross section occurs at high centre of mass energy for the process e+e-  $\rightarrow$  H0A0 as compared to that of e+e-  $\rightarrow$  h0A0.

Primary author: Mr OUHAMMOU, Mohamed (sultan moulay slimane university)
Co-authors: Mr OUALI, Moha; Prof. TAJ, Souad; Prof. MANAUT, Bouzid; Prof. BENBRIK, Rachid
Presenter: Mr OUHAMMOU, Mohamed (sultan moulay slimane university)
Session Classification: Parallel Session II