## On 6D N=(1,0) Supergravity

#### R. Sammani, Y. Boujakhrout, R. Ahl Laamara

#### LPHE-MS, Science faculty, Mohammed V University in Rabat, Morocco.

#### Orator: R. Sammani , March 22, 2022



- Motivations and Introduction
- The anomaly constraints
- The moduli space constraints
- The BPS string constraints
- Geometric constraints
- Summary and Future directions

- Goal : Build a quantum gravity theory in 4D.
- **Problem** : We have a very limited knowledge of the type of constraints imposed by UV completion in 4D.
- Current Works : Find another space to know more about types of constraints.

Why 6D N=(1,0) supergravity space?

- Successful Landscape analysis : Strong constraints.
- **F-theory realizations** : 6D supergravity models are naturally realized within F-theoty compactified on three-folds.
- 8 supersymmetries : We want to expand the work for 8 supersymmetries but 4D N = 2.

### Introduction

Consistent quantum field theories when coupled to gravity can be categorized as :



what are those constraints and conditions that could make a theory consistent or not  $? \end{tabular}$ 

R. Sammani, Y. Boujakhrout, R. Ahl Laamara

On 6D N=(1,0) Supergravity





To investigate whether the theory has anomalies or not, we need to express the anomaly polynomial.

Anomalies can be cancelled by the GreenSchwarz-Sagnotti mechanism if the anomaly polynomial  $I_8$  factorizes as :  $I_8(R, F) = \frac{1}{2}\Omega_{\alpha\beta}X_4^{\alpha}X_4^{\beta}$  $X_4^{\alpha} = \frac{1}{2}a^{\alpha}trR^2 + \Sigma b_i^{\alpha}\frac{2}{\lambda_i}trF_i^2$ 

So our goal is to express the anomaly polynomial in 6d dimensions and to factorize it so it can be cancelled.



#### Anomalies

• Gauge Anomaly This type of anomaly in 6d is due to the antichiral spin fermion in the hypermultiplet and to the chiral spin fermion in the vector multiplet :

 $I_8^{gauge, \frac{1}{2}} = -\frac{1}{24(2\pi)^3} Tr_R F^4 + \frac{1}{24(2\pi)^3} \sum n_R^i Tr_R F^4 + \frac{1}{4(2\pi)^3} \sum n_R^i n_S^i Tr_R F^2 Tr_S F^2$ R/S are representations of the considered group.  $n_R^i$  is the number of the hypermultiplet in the representation R.

Gravitational Anomaly They are contributions from the gravitino, the chiral and the anti-chiral fermions in the vector, tensor and hype rmultiplets. Adding also the contribution from the supergravity 3-form field : The total gravitational anomaly is :

 $I_8^{\text{grav}} = \frac{1}{24(2\pi)^3} (\frac{1}{240} (H - V + 29T - 273) tr R^4 + \frac{1}{192} (H - V - 7T + 51) (tr R^2)^2)$ 

Mixed Anomaly Due to the chiral and the anti chiral spin fermions in vector and hyper multiplets :

$${}_{8}^{\text{mixed},\frac{1}{2}} = \frac{1}{(2\pi)^{3}} \left(\frac{1}{96} tr R^{2} Tr_{R} F^{2} - \frac{1}{96} tr R^{2} \Sigma n_{R}^{i} Tr_{R} F^{2}\right)$$

We sum all the anomalies above to get :

$$(2\pi)^{3}I_{8} = \frac{1}{5760}(H - V + 244)trR^{4} + \frac{1}{4608}(H - V + 44)(trR^{2})^{2} + \frac{1}{96}trR^{2}(Tr_{R}F^{2} - \Sigma n_{R}^{i}Tr_{R}F^{2}) - \frac{1}{24}Tr_{R}F^{4} + \frac{1}{24}\Sigma n_{R}^{i}Tr_{R}F^{4} + \frac{1}{4}\Sigma n_{RS}^{ij}Tr_{R}F^{2}Tr_{S}F^{2}$$

By taking T=1, one tensor multiplet.

However, the term  $trR^4$  cannot be factorized in the said way, thus the term associated with it needs to vanish : H-V+244=0

To transform the trace from R representation to the fundamental representation, we use :

$$Tr_R F^2 = A_R tr F^2, Tr_R F^4 = B_R tr F^4 + C_R (tr F^2)^2$$

Finally, we have :

$$I_8 \propto \left\{ (trR^2)^2 + \frac{1}{6}trR^2 (A_R - \sum n_R^i A_R^i) trF^2 - \frac{2}{3} (B_R - \sum n_R^i B_R^i) trF^4 - \frac{2}{3} (C_R - \sum n_R^i C_R^i) (trF^2)^2 + 4\sum n_{RS}^{ij} tr_R F^2 tr_S F^2 A_R A_S \right\}$$

The goal is to write is as :

$$I_8(R,F) = \frac{1}{2}\Omega_{\alpha\beta}X_4^{\alpha}X_4^{\beta}$$
 where  $X_4^{\alpha} = \frac{1}{2}a^{\alpha}trR^2 + \Sigma b_i^{\alpha}\frac{2}{\lambda_i}trF_i^2$ 

Thus :

$$I_{8}(R,F) = \frac{1}{2} \left\{ \frac{a^{\alpha}a^{\beta}}{4} (trR^{2})^{2} - 2 \frac{\sum a^{\alpha}b_{j}^{\beta}}{\lambda_{j}} trR^{2} trF^{2} + 4 \frac{\sum b_{i}^{\alpha}\sum b_{j}^{\beta}}{\lambda_{i}\lambda_{j}} trF_{i}^{2} trF_{j}^{2} \right\}$$

### The anomaly constraints

The anomaly factorization conditions for gravitational, gauge and mixed anomalies are summarized as follows :

$$\begin{array}{l} \mathbb{R}^{4} \longrightarrow H - V + 244 = 0 = 273 - 29T \\ \mathbb{F}^{4} \longrightarrow \mathbb{B}_{R} = \Sigma n_{R}^{i} \mathbb{B}_{R}^{i} \\ (\mathbb{R}^{2})^{2} \longrightarrow a a = 8 = 9 - T \\ \mathbb{F}^{2} \mathbb{R}^{2} \longrightarrow a b_{j} / \lambda_{j} = \frac{1}{6} (A_{R} - \Sigma n_{R}^{i} A_{R}^{i}) \\ (\mathbb{F}^{2})^{2} \longrightarrow b_{i} b_{j} / \lambda_{i}^{2} = \frac{1}{3} (\Sigma n_{R}^{i} C_{R}^{i} - C_{R}) \\ (\mathbb{F}_{i}^{2}) (\mathbb{F}_{j}^{2}) \longrightarrow b_{i} b_{j} / \lambda_{i} \lambda_{j} = (\Sigma n_{RS}^{ij} A_{R} A_{S}) \end{array}$$

#### The anomaly constraints

The anomaly factorization conditions for gravitational, gauge and mixed anomalies are summarized as follows :

$$\begin{array}{l} \mathbb{R}^{4} \longrightarrow H - V + 244 = 0 = 273 - 29T \\ \mathbb{F}^{4} \longrightarrow \mathbb{B}_{R} = \Sigma n_{R}^{i} B_{R}^{i} \\ (\mathbb{R}^{2})^{2} \longrightarrow a a = 8 = 9 - T \\ \mathbb{F}^{2} \mathbb{R}^{2} \longrightarrow a b_{j} / \lambda_{j} = \frac{1}{6} (A_{R} - \Sigma n_{R}^{i} A_{R}^{i}) \\ (\mathbb{F}^{2})^{2} \longrightarrow b_{i} b_{j} / \lambda_{i}^{2} = \frac{1}{3} (\Sigma n_{R}^{i} C_{R}^{i} - C_{R}) \\ (\mathbb{F}_{i}^{2}) (\mathbb{F}_{j}^{2}) \longrightarrow b_{i} b_{j} / \lambda_{i} \lambda_{j} = (\Sigma n_{RS}^{ij} A_{R} A_{S}) \end{array}$$

The vectors,  $a^{\alpha}$ ,  $b_i^{\alpha}$  are constrained to have integer inner products a \* a,  $a * b_j$ ,  $b_i * b_i$ . We call this the anomaly lattice.

$$\Lambda = \begin{pmatrix} a^2 & -a \cdot b_1 - a \cdot b_2 \cdots \\ -a \cdot b_1 & b_1^2 & b_1 \cdot b_2 \cdots \\ -a \cdot b_2 & b_1 \cdot b_2 & b_2^2 \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

R. Sammani, Y. Boujakhrout, R. Ahl Laamara

#### Also...

The 6D theory has gravity/gauge dyonic strings with charges a,  $b_i$ . Those charges span the anomaly lattice which is contained in the full string lattice of the 6d theory. The anomaly lattice is required to have a unimodular embedding into a self-dual lattice and this fact provides a constraint on possible theories.

How does this work?

# Example : 6D supergravity theory with $\mathsf{T}=1,$ gauge group $\mathsf{G}{=}\mathsf{SU}(4)$

The matter content is :



Now we need to write the anomaly lattice.

$$a.b = \frac{\lambda}{6} (A_{adj} - \sum n_R A_R) = \frac{1}{6} (8 - (8 + 40 + 1 + 10 + 2) = -10$$

$$a.a = 9 - T = 9 - 1 = 8$$

$$A = \begin{pmatrix} a.a \\ -a.b \\ b.b = \frac{\lambda^2}{3} (\sum n_R C_R - C_{adj}) = \frac{1}{3} (6 + 40 + 0 + 10 + 3 - 6) = 10$$

$$b.b = \frac{\lambda^2}{3} (\sum n_R C_R - C_{adj}) = \frac{1}{3} (6 + 40 + 0 + 10 + 3 - 6) = 10$$

$$b.b = \frac{\lambda^2}{3} (\sum n_R C_R - C_{adj}) = \frac{1}{3} (6 + 40 + 0 + 10 + 3 - 6) = 10$$

$$b.b = \frac{\lambda^2}{3} (\sum n_R C_R - C_{adj}) = \frac{1}{3} (2 + 40 + 0 + 10 + 3 - 6) = 10$$

# Example : 6D supergravity theory with $\mathsf{T}=1,$ gauge group $\mathsf{G}{=}\mathsf{SU}(4)$

 $\wedge = \begin{pmatrix} 8 & 10 \\ 10 & 10 \end{pmatrix}$ 

The lattice above needs to have a perfect square determinant. Since the determinant is -20, not a perfect square than the lattice does not admit an embedding into any unimodular lattice.

The moduli space of the 6d (1,0) supergravity locally takes the form SO(1,T)/SO(T) which is parametrized by the a vector j with positive norm j.j > 0 representing the positivity of the metric on the moduli space.

The next set of constraints are required for the proper sign of the gauge kinetic term and gravitational term.

Actually, The gauge kinetic term is proportional to  $-j.b_i tr F^2$ . Thus we require that that  $j.b_i > 0$ .

The gravitational term is proportional to  $j.atrR^2$ , we require consequently that j.a < 0.

The moduli space of the 6d (1,0) supergravity locally takes the form SO(1,T)/SO(T) which is parametrized by the a vector j with positive norm j.j > 0 representing the positivity of the metric on the moduli space.

The next set of constraints are required for the proper sign of the gauge kinetic term and gravitational term.

Actually, The gauge kinetic term is proportional to  $-j.b_i tr F^2$ . Thus we require that that  $j.b_i > 0$ .

The gravitational term is proportional to  $j.atrR^2$ , we require consequently that j.a < 0.

For example, if a theory has :  $b = \lambda a, \lambda > 0$ , it will be inconsistent because we can't find any a, b that verify simultaneously j.a < 0 and j.b > 0.

### BPS string consideration

The existence of the two-form fields  $B_2$  implies the existence of string sources in accordance with the hypothesis that the spectrum of a gravitational theory needs to be complete.

The worldsheet theory on those strings gives rise to 2d SCFT at low energy with charge Q. Next, we need to express the anomaly inflow from 6d to 2d :

$$I_4 = -I_4^{inflow} = \Omega_{lphaeta} Q^lpha (X_4^lpha + rac{1}{2} Q^eta (c(l) - c(R)))$$

since :

$$X_4^{\alpha} = \frac{1}{2}a^{\alpha}trR^2 + \frac{1}{4}\Sigma b_i^{\alpha}trF_i^2$$

then :

$$I_4 = \Omega_{\alpha\beta} Q^{\alpha} (\frac{1}{2} a^{\alpha} tr R^2 + \frac{1}{4} \Sigma b_i^{\alpha} tr F_i^2 + \frac{1}{2} Q^{\beta} (c(l) - c(R))$$

we have :

$$trR^2 = \frac{-1}{2}p(T) + c(I) + c(R)$$

Then :

$$I_4 = \frac{-Q.a}{4} p(T) + \frac{1}{4} \Sigma Q.b_i^{\alpha} tr F_i^2 - \frac{Q.Q-Q.a}{2} c(R) + \frac{Q.Q+Q.a}{2} c(I)$$

R. Sammani, Y. Boujakhrout, R. Ahl Laamara

$$I_4 = \frac{-Q.a}{4}p(T) + \frac{1}{4}\Sigma Q.b_i^{\alpha}trF_i^2 - \frac{Q.Q-Q.a}{2}c(R) + \frac{Q.Q+Q.a}{2}c(I)$$

By eliminating the contribution from the center of mass degrees of freedom :

$$I_4 \longrightarrow I_4 - \left(-\frac{1}{12}p(T) - c(I)\right)$$

Which gives :



Finally, since the central charges need to be positive :

$$Q.Q \geq -1, k_l = Q.Q + Q.a + 2 \geq 0, k_i = Q.b_i \geq 0$$

A worldsheet theory has non-negative tension only if  $Q \cdot j \ge 0$ . Only strings with  $Q \cdot j \ge 0$  embedded in 6d supergravity theories give rise to unitary 2d SCFTs.

The second condition that rises from SCFT : The central charge of group G is,

$$c_G = \frac{k.dimG}{k+h}$$

where k is the  $k_i$  defined by Q.b previously and h is the dual Coxeter number of group G, for SU(N) is equal to N.

The SCFT states that, the current algebra for group G is on the left-moving sector. This tells us that the sum of the group central charges is less than  $c_L$ . Let's take an example

The example is the 6d supergravity with T = 1 and SU(N) gauge group with :

$$\Omega = (1, -1), a = (-3, 1), b = (0, -1), j = (n, 1)$$

Let's derive the consistency conditions for  $Q = (q_1, q_2)$ .

$$egin{aligned} Q.Q &\geq -1 \implies q_1^2 - q_2^2 \geq -1 \ Q.Q + Q.a + 2 \geq 0 \implies q_1^2 - q_2^2 - 3q_1 - q_2 \geq -2 \ k_i &= Q.b_i = q_2 \geq 0 \ j.Q &= nq_1 - q_2 \geq 0 \ \sum c_{Gi} &= \sum rac{k.dimG}{k+h} \leq c_L = 3Q.Q - 9Q.a + 2 \ &\implies rac{q_2(N^2-1)}{q_2+N} \leq 3(q_1^2 - q_2^2) + 9(3q_1 + q_2) + 2 \end{aligned}$$

6D N = (1, 0) supergravity models are 'naturally' realized in the context of F-theory : a large class of 6D supergravity theories are realized by F-theoty compactified on three-folds.



Up till now, our theory has three major parameters : a, b and j. So any requirement of f-theory needs to evolve around them.

Up till now, our theory has three major parameters : a, b and j. So any requirement of f-theory needs to evolve around them.

One of F-theory constraints on low-energy supergravity is the Kodaira condition :

$$-12K_B = \sum \nu_i \epsilon_i + Y$$

with,

J.Y > 0

Where  $K_B$ , is a canonical divisor.  $\epsilon_i$ , are effective, irreducible curves. Y, is a residual divisor and J is a Kahler class .

Up till now, our theory has three major parameters : a, b and j. So any requirement of f-theory needs to evolve around them.

One of F-theory constraints on low-energy supergravity is the Kodaira condition :

$$-12K_B = \sum \nu_i \epsilon_i + Y$$

with,

J.Y > 0

Where  $K_B$ , is a canonical divisor.  $\epsilon_i$ , are effective, irreducible curves. *Y*, is a residual divisor and *J* is a Kahler class .

By constructing the following mapping from 6d sugra to f-theory :

$$\begin{array}{c} a \Longrightarrow K_B \\ b \Longrightarrow \epsilon_i \\ j \Longrightarrow J \end{array}$$

The Kodaira condition becomes :

$$j(-12a - \sum 
u_i b_i) \geq 0$$

Additional constraints on both a and  $b_i$  could be formulated based on the previous map. Since :

$$-K_B = (3, -1, -1..., -1)$$

Then, we have :

 $a = (-3, 1^T)$ 

Also :

Then :

 $\epsilon \ast \epsilon' < \mathsf{0}$ 

 $b^{2} < 0$ 

## Summary Future directions

