



Contribution ID: 76

Type: **not specified**

## Charged Higgs boson production via $pp \rightarrow H^\pm b j$ at the LHC

*Wednesday, 23 March 2022 15:00 (15 minutes)*

The charged Higgs searches can be served to probe new physics at the LHC. In this study, we focus on the associated production of the charged Higgs boson with the bottom-quark and jet in 2HDM-type-I as a promising mode for a light  $H^\pm$ , i.e.  $m_{H^\pm} < m_t$ . We consider both situations where  $h(H)$  are the SM-like Higgs boson discovered with a mass near 125 GeV and investigate their bosonic decays, such as  $H^\pm \rightarrow W^\pm h$  and/or  $H^\pm \rightarrow W^\pm A$ . We explore the possible signals at the LHC taking into account the theoretical and experimental constraints, as a result, we find that, over a substantial region of the 2HDM-I parameter space, the Signal  $qbW + 2b/2\tau/2\gamma$  could serve as a promising and alternative signal that might serve to discover the  $H^\pm$  states at the LHC.

**Primary authors:** OUCHEMHOU, Mohamed (Cadi Ayyad University, Marrakech, Morocco.); Prof. ARHRIB, Abdesalam (Abdelmalek Essaadi University, Faculty of Sciences and techniques, Tanger, Morocco.); Prof. BENBRIK, Rachid (Laboratoire de Physique Fondamentale et Applique Safi, Faculté Polydisciplinaire de Safi, Sidi Bouzid, B.P. 4162, Safi, Morocco.); Mr KRAB, Mohamed (Research Laboratory in Physics and Engineering Sciences, Modern and Applied Physics Team, Polidisciplinary Faculty, Beni Mellal, 23000, Morocco.)

**Presenter:** OUCHEMHOU, Mohamed (Cadi Ayyad University, Marrakech, Morocco.)

**Session Classification:** Parallel Session V, Collider - Experiment