

Testing indirect experimental methods for constraining the $^{193,194}\text{Ir}(n,\gamma)$ cross sections

S. P. E. Magagula^{1,2}, A. C. Larsen³, and M. Wiedeking^{1,2}

¹SSC laboratory, iThemba LABS, P.O. Box 722, Somerset West
7129, South Africa.,

²School of Physics, University of the Witwatersrand, Johannesburg
2050, South Africa.,

³Department of Physics, University of Oslo, N-0316 Oslo, Norway,

July 2023

Abstract

As much as nucleosynthesis or element formation is concerned, almost all the nuclei heavier than iron have been made in part by the slow neutron capture and the rapid neutron capture processes ($\approx 50\%$ each), respectively known as the s- and r- processes [1]

The neutron capture reactions $^{192}\text{Ir}(n,\gamma)^{193}\text{Ir}$ and $^{193}\text{Ir}(n,\gamma)^{194}\text{Ir}$ are indirectly studied by analysing data obtained from the Oslo Cyclotron Laboratory (OCL). These data will allow for the study of $^{193,194}\text{Ir}$ isotopes, from the $^{192}\text{Os}(\alpha,t\gamma)$ and $^{192}\text{Os}(\alpha,d\gamma)$ reactions, respectively. The $^{193}\text{Ir}(n,\gamma)^{194}\text{Ir}$ cross sections which will be constrained by our measurement will provide a comparison to existing (n,γ) measurement data [2].

In addition, the $^{192}\text{Ir}(n,\gamma)^{193}\text{Ir}$ reaction is a branching point in the s-process making it very interesting, but it is challenging to measure the (n,γ) cross section directly since ^{192}Ir is unstable. Therefore the OCL data may provide very valuable information on the $^{192}\text{Ir}(n,\gamma)^{193}\text{Ir}$ cross section by indirectly constraining it with the experimental nuclear level density (NLD) and γ -strength function (γSF).

An array of Sodium Iodine (NaI)Tl detectors, called CACTUS, detected γ -rays and the silicon particle telescope array, called SiRi, was used to detect charged particles in coincidence. The NLDs and γSF s are being extracted below the neutron separation energy, S_n , using the Oslo Method [3]. Furthermore, the NLDs and γSF s will be used as inputs in the open-source code called TALYS to calculate cross-sections of $^{193,194}\text{Ir}$. I will provide preliminary results of the measured NLDs and γSF s from the $^{192}\text{Os}(\alpha,d\gamma)^{194}\text{Ir}$ reaction which will be used as inputs in the code TALYS to calculate cross-sections of $^{193,194}\text{Ir}$.

- [1] Arnould, M., Goriely, S., and Takahashi, K. (2007). *Physics Reports*, 450(4-6), 97-213.
- [2] Zerkin, V. V., and Pritychenko, B. (2018). The experimental nuclear reaction data (EXFOR) 888, 31-43.
- [3] Schiller, A., Bergholt, L., Guttormsen, M., Melby, E., Rekstad, J., and Siem, S. (2000). *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 447(3), 498-511.

This work is based on the research supported in part by the National Research Foundation of South Africa (Grant Number: 118846) and the International Atomic Energy Agency under Research Contract 20454.