



Contribution ID: 321

Type: Oral

Sub-barrier Fusion Excitation Functions of Heavy-Ion Systems

Saturday, 2 December 2023 12:20 (15 minutes)

One of the yet unsettled problems in heavy-ion fusion near and below the barrier is the relative influence of nucleon transfer channels and couplings to collective modes on the cross sections. We recall two relevant papers [1] where that influence, and the moments of fusion-barrier distributions were investigated.

In this contribution, we present a new analysis of several systems, based on the combined observation of the energy-weighted excitation functions $E\sigma$ in relation to their first energy derivatives $d(E\sigma)/dE$ (slopes). That derivative is proportional to the s-wave transmission coefficient and to the square of the barrier radius. This representation helps our understanding of the situation.

The two-dimensional plot of $d(E\sigma)/dE$ vs $E\sigma$ for $^{48}\text{Ca}, ^{36}\text{S} + ^{48}\text{Ca}$, obtained from Refs. [2, 3] is interesting. In this type of plot trivial Coulomb barrier differences between the two systems are eliminated to a large extent. The colliding nuclei are closed-shell or magic, and, at sub-barrier energies, the two data sets are completely overlapping. Indeed the Wong formula [4] implies that the slope and the excitation function are proportional to each other for all cross sections in that energy range. The proportionality constant is $2\pi/\hbar^2\omega$, i.e. inversely proportional to the second radial derivative of the barrier approximated by a parabola. In the cited example the overlap of the two data sets implies that the two barriers have approximately the same width.

Other cases behave differently, like $^{40}\text{Ca} + ^{96}\text{Zr}$ [5] and $^{58}\text{Ni} + ^{64}\text{Ni}$ [6], where neutron transfer couplings are dominant, compared to $^{40}\text{Ca} + ^{90}\text{Zr}$ [7] and $^{64}\text{Ni} + ^{64}\text{Ni}$ [8], respectively. In either case, the system where transfer couplings are dominant, lies below the other case, meaning that the effective one-dimensional barrier is thinner. This mimics a wider barrier distribution produced by couplings, extending to lower energies and leading to a cross section enhancement vs energy, as observed. A full systematics will be shown in the talk, with more detailed and quantitative considerations for the various cases.

- [1] C. L. Jiang et al., Phys. Rev. C 89 (2014) 051603(R); K. E. Rehm et al., Phys. Rev. C 94 (2016) 044612
- [2] A.M. Stefanini, G. Montagnoli et al., Phys. Lett. B 679 (2009) 95
- [3] A.M. Stefanini, G. Montagnoli et al., Phys. Rev. C 78 (2008) 044607
- [4] C. Y. Wong, Phys. Rev. Lett. 31 (1973) 766
- [5] A.M. Stefanini, G. Montagnoli et al., Phys. Lett. B 728 (2014) 639
- [6] A.M. Stefanini, G. Montagnoli, M. Del Fabbro et al., Phys. Rev. C 100 (2019) 044619
- [7] H. Timmers et al., Nucl. Phys. A 633 (1998) 421
- [8] C. L. Jiang et al. Phys. Rev. Lett. 93 (2004) 012701

Attendance Type

Remote

Primary authors: STEFANINI, Alberto M. (INFN - LNL); MONTAGNOLI, Giovanna (Università di Padova); DEL FABBRO, Mirco (Dipartimento di Fisica e Scienze della Terra Univ. di Ferrara and INFN Padova); Dr CORRADI, Lorenzo (INFN - LNL); FIORETTO, Enrico (INFN - LN L); SZILNER, Suzana (dr)

Presenter: STEFANINI, Alberto M. (INFN - LNL)

Session Classification: Session 10

Track Classification: Nuclear Structure, Reactions and Dynamics