



Contribution ID: 16

Type: Oral

Microscopic Analysis of Elastic Scattering of One-Proton Halo Nucleus ^{17}F on Different Mass Targets

An analysis of cross sections of elastic scattering of ^{17}F on ^{12}C , ^{14}N , ^{58}Ni , and ^{208}Pb nuclei at energy 170 MeV and on ^{208}Pb at various energies is carried out by using the microscopic optical potentials (OPs) [1]. The proton and neutron density distributions of the exotic nucleus ^{17}F are computed in the framework of microscopic models. The real part of the OP is calculated by a corresponding folding procedure accounting for the antisymmetrization effects, while the imaginary part is obtained on the base of the high-energy approximation [2]. In the hybrid model of the optical potential developed and explored in our previous works [3,4] the only free parameters are the depths of the real and imaginary parts of the OPs obtained by fitting the experimental data. A good agreement of the theoretical results with the available experimental data is achieved pointing out clearly to a peripheral character of the scattering.

References

- [1] M.K. Gaidarov, K.V. Lukyanov, E.V. Zemlyanaya, V.K. Lukyanov, D.N. Kadrev, and A.N. Antonov, Phys. Part. Nucl. 54, 500 (2023).
- [2] V.K. Lukyanov, E.V. Zemlyanaya, and K. V. Lukyanov, Phys. At. Nucl. 69, 240 (2006); JINR Preprint P4-2004-115, Dubna, 2004.
- [3] K.V. Lukyanov et al., Eur. Phys. J. A 33, 389 (2007).
- [4] V.K. Lukyanov et al., Phys. Rev. C 80, 024609 (2009); Phys. Rev. C 82, 024604 (2010); Phys. Rev. C 88, 034612 (2013); Phys. Rev. C 91, 034606 (2015); Eur. Phys. J. A 53, 31 (2017); Phys. Rev. C 100, 034602 (2019).

Notes

Primary authors: Prof. GAIDAROV, M.K. (INRNE-BAS); Dr LUKYANOV, K.V. (JINR, Dubn); Prof. ZEMLYANAYA, E.V. (JINR, Dubna, Russia); Prof. LUKYANOV, V.K. (JINR, Dubna, Russia); Dr KADREV, D.N. (INRNE-BAS); Prof. ANTONOV, A.N. (INRNE-BAS)

Presenter: Prof. GAIDAROV, M.K. (INRNE-BAS)