



Contribution ID: 87

Type: Oral

In a search for the true identity of low-lying positive parity states in the $A \sim 160$ mass region.

For many years, the first excited 0^+ bands of well-deformed nuclei were understood as “ β -vibrations”, following the seminal works of Bohr and Mottelson [1, 2]. Nevertheless, over the years it has become apparent that low-lying 0^+ bands could also arise due to other effects such as shape-coexistence, and quadrupole pairing [3, 4, 5].

In this work, I will present a comprehensive dataset comprising of low-lying positive parity bands (β and γ bands) in even-even isotopes with $N \sim 88$ to 92 and proton numbers $Z \sim 62$ (Sm) to 70 (Yb). In order to produce a complete and definitive microscopic picture of the so-called β bands as well as other related low-lying positive parity excitations, an extensive systematic review is performed for these nuclei. The implication of the findings on the interpretation of the first excited 0^+ states and other related low-lying positive-parity excitations is there from discussed.

Reference

- [1] A. Bohr, Mat. Fys. Medd. Dan. Vid. Selsk. 26, 14 (1952).
- [2] A. Bohr and B. R. Mottelson, Mat. Fys. Medd. Dan. Vid. Selsk. 27, 16 (1953).
- [3] K. Heyde and J. L. Wood, Reviews of Modern Physics 83, 1467 (2011).
- [4] J.F. Sharpey-Schafer et al., Eur. J. Phys. A 47, 5 (2011), Eur. J. Phys. A 47, 6 (2011).
- [5] P.E. Garrett, J Phys. G: Nucl. Part. Phys. 27, R1 (2001).

Notes

Primary authors: Dr MAJOLA, Siyabonga (University of Johannesburg); MAJOLA, Siyabonga (UJ)

Presenter: Dr MAJOLA, Siyabonga (University of Johannesburg)