



Contribution ID: 27

Type: Oral

## Study of the spatial dimensions of ${}^6\text{Li}$ nuclear states by measuring differential cross sections for the ${}^{10}\text{B}({}^7\text{Li}, {}^6\text{Li}){}^{11}\text{B}$ reaction

The angular distributions of the 1-n transfer reaction  ${}^{10}\text{B}({}^7\text{Li}, {}^6\text{Li}){}^{11}\text{B}$ , as well as the elastic scattering of  ${}^7\text{Li}$ , were measured at  $E_{\text{lab}} = 58$  MeV. Experiment was done using U-400 accelerator beam of the FLNR JINR, Dubna.

The attention was paid to ground state (g.s.) and excited ( $J=0^+$ ,  $T=1$ ,  $E=3.56$  MeV) state of  ${}^6\text{Li}$  which is an isobar analogue state (IAS) of 2n-halo nucleus  ${}^6\text{He}$ .

Angular distribution for 1-step direct reaction  ${}^{10}\text{B}({}^7\text{Li}, {}^6\text{Li}){}^{11}\text{B}$  with excitation of the 3.56 MeV state ( ${}^6\text{Li}$ ) is present for the first time.

The DWBA analysis of the differential cross section of the  ${}^{10}\text{B}({}^7\text{Li}, {}^6\text{Li}){}^{11}\text{B}$  the  ${}^6\text{Li}_{\text{g.s.}}$  and  ${}^6\text{Li}(3.56$  MeV) transition was performed. The optical model potentials were obtained by fitting of measured elastic scattering data and evaluating parameters for the output reaction channels. Phenomenological approaching based on solving an approximate equation for the reaction form factor was used to determine its radial dependence and empirical values of asymptotic normalization coefficient (ANC). Obtained values of ANC's for the  ${}^6\text{Li}_{\text{g.s.}}$  and  ${}^6\text{Li}(3.56$  MeV) states are similar to literature one. Comparison of the radial dependences of form factors shows that the wave function of the  ${}^6\text{Li}$  nucleus in excited ( $J=0^+$ ,  $T=1$ ,  $E=3.56$  MeV) state has increased spatial dimension compared to the ground state, and in both cases some larger spatial size than the ground states of the  ${}^{11}\text{B}$  and  ${}^{10}\text{B}$ . Within the framework of our analysis, we can confirm that the radius of the  ${}^6\text{Li}$  nucleus in the 3.56 MeV state is larger than in the ground state.

This result, obtained within the framework of the ANS analysis, is an argument in favor of the existence of a halo in the  ${}^6\text{Li}(3.56$  MeV) state, while the question of a halo in  ${}^6\text{Li}_{\text{g.s.}}$  still leaves open.

### Notes

**Primary author:** Dr SOBOLEV, Yuri (Joint Institute for Nuclear Research)

**Co-authors:** Mr STUKALOV, Sergei; Prof. PENIONZHKEVICH, Yuri; Dr BURTEBAUEV, Nassurulla (Institute of Nuclear Physics of the Ministry of Energy of the Republic of Kazakhstan); Dr GONCHAROV, Sergey (M.V. Lomonosov Moscow State University); Dr GUROV, Yuri (Joint Institute for Nuclear Research); Dr DANILOV, Andrey (National Research Centre Kurchatov Institute); Dr DEMYANOVA, Alla (National Research Centre Kurchatov Institute); Dr NASSURLLA, Maulen (Institute of Nuclear Physics of the Ministry of Energy of the Republic of Kazakhstan.); Mr STARASTSIN, Viktor (National Research Centre Kurchatov Institute); Mr SHAKHOV, Alexey (Joint Institute for Nuclear Research)

**Presenter:** Dr SOBOLEV, Yuri (Joint Institute for Nuclear Research)