



Contribution ID: 57

Type: Oral

Two-neutron-transfer to ^{178}Yb and Population of $^{178m2}\text{Hf}$ via Incomplete Fusion

The DIAMANT light-charged-particle detector from ATOMKI was coupled with the AFRODITE gamma-ray spectrometer at iThemba LABS in a collaboration enabled by a bilateral agreement between the governments of South Africa and Hungary. This facilitated the study of incomplete fusion reactions in the bombardment of a Ytterbium-176 target with a beam of 50 MeV Lithium-7 ions. The beam was generated as a collaborative effort between ion source experts at iThemba LABS and the Flerov Laboratory for Nuclear Reactions (FLNR) of the Joint Institute for Nuclear Reactions (JINR), Dubna.

Particle-Identification (PID) spectra from DIAMANT generated from the ATOMKI custom-built VXI electronics clearly show the detection of protons, tritons and alpha particles, which, when gated on, allowed the selection of gamma-ray coincidences detected with AFRODITE when the respective complementary Helium-6, Helium-4 (α) and triton fragments fused with the target.

Analysis of the charged-particle-selected gamma-ray coincidence data enabled the identification of Hafnium-180 in the proton-gated E_γ - E_γ correlation matrix, as well as Hafnium-178, including the band based on the $T_{1/2} = 31\text{a}$, $K^\pi = 16^+$ four-quasiparticle state. Hafnium-178 is also evident in the triton-gated matrix, which suggests that this nucleus is populated via two incomplete fusion channels, this one in which the fused fragment is Helium-4, and the other in which a Helium-6 neutron-rich fragment fuses with the Ytterbium-176 target.

The relative contribution from the $(^7\text{Li}, p4n)$ fusion evaporation channel is unclear, but there is other evidence for Helium-6-induced reactions in the population of neutron-rich Ytterbium-178 whereby two neutrons have been transferred to the target. The ground-state band of Ytterbium-178 can be clearly observed in both the proton-gated and alpha-gated matrices, which supports the assignment of the $^{176}\text{Yb}(^7\text{Li}, \alpha p)^{178}\text{Yb}$ reaction. The deuteron yield is comparatively weak which has hampered the unambiguous confirmation of the $^{176}\text{Yb}(^7\text{Li}, \alpha d)^{177}\text{Yb}$ reaction.

The comparatively strong population of Hafnium-178 via the two reaction channels discussed above has allowed the population ratio $I_\gamma(\text{proton-gated})/I_\gamma(\text{triton-gated})$ of the ground-state, two-quasiparticle $K^\pi = 8^-$ and four-quasiparticle $K^\pi = 14^-$ and $K^\pi = 16^+$ bands to be extracted as function of spin. There is evidence for a marked increase in relative population of the $K^\pi = 16^+$ band when compared to the other lower-spin band structures.

Notes

Primary authors: Dr MULLINS, Simon (Botswana International University of Science & Technology); Dr MAQABUKA, Bongani (University of Johannesburg); Dr BARK, Robert (iThemba LABS); Dr BOGOLOMOV, Sergei (JINR); Prof. CONNELL, Simon (University of Johannesburg); Dr ANDREI, Efremov (JINR); Dr KUTI, Istvan (ATOMKI); Dr LAWRIE, Elena (iThemba LABS); Prof. LAWRIE, Kobus (University of the Western Cape); Dr MAJOLA, Siyabonga (University of Johannesburg); Dr MOLNAR, Jozsef (ATOMKI); Mr MURRAY, Sean (University of Cape Town); Dr NYAKO, Barna (ATOMKI); Dr PAPKA, Paul; Dr THOMAE, Rainer

Presenter: Dr MULLINS, Simon (Botswana International University of Science & Technology)