

Nuclear structure made clear by Gamow-Teller transitions

Gamow-Teller (GT) transitions are caused by the σ ; τ ; operator. Therefore, they reflect the uniqueness of nuclei that consist of two types of Fermions “protons and neutrons” with “spin-degree” of freedom. In addition, GT transitions are the most common nuclear weak transitions that are active all over the Universe. Most direct information on GT transition strength $B(GT)$ can be derived from beta-decay measurements, but they cannot access highly excited states. In the 1980s, it was found that p,n charge-exchange (CE) reactions at intermediate energies ($E_{p,n} > 100$ MeV) and 0-degrees are sensitive to the σ ; τ ; response of nuclei. Thus, they became the break through against the Q -value limitation in the beta-decay study. In the late 1990s, ($^3\text{He}, t$), CE reaction at 0-degrees was introduced, in which a magnetic spectrometer was used for the analysis of tritons. By the realization of beam matching conditions for the high-dispersive beam line WS course and the Grand Raiden spectrometer, high energy-resolutions of 30 keV or even better were achieved at the incoming ^3He energy of 140 MeV/nucleon. The overview of the study of GT transitions at RCNP, Osaka [1] are presented for p -shell, sd -shell, pf -shell, and also for heavier mass A nuclei. We see that they are the transitions with full of personality.

[1] Y. Fujita, B. Rubio and W. Gelletly, Prog. in Part. and Nucl. Phys. 66, 549 (2011).

Primary author: Prof. FUJITA, Yoshitaka (RCNP & Department of Physics, Osaka Univeristy)

Co-author: Dr FUJITA, Hirohiko (RCNP & Department of Physics, Osaka Univeristy)

Presenter: Prof. FUJITA, Yoshitaka (RCNP & Department of Physics, Osaka Univeristy)