


An effective model description of QCD thermodynamics¹

Ludwik Turko

University Wrocław, Poland

On Discovery Physics at the LHC
Kruger 2012

December 3 - 7, 2012



¹based on paper by D. Blaschke, D. Prorok, L.T. and J. Berdermann - Acta Phys. Polon. Supp. **5**, 485 (2012) *and in further progress*




To achieve

A compound and consistent effective model reproducing:


- The equation of state of hadronic matter as obtained in lattice QCD simulations.
- Basic physical characteristics of processes encountered in the dense hadronic matter:
 - Modified in medium properties of hadrons as different from those in the vacuum.
 - Modification of the notion of the mass shell.
 - Dissolving of hadrons into quarks and gluons phase,
 - Physical processes present in the full QCD treatment.



Theoretical laboratory of QCD



The energy density normalized by T^4 as a function of the temperature on $N_t=6,8$ and 10 lattices.



The pressure normalized by T^4 as a function of the temperature on $N_t=6,8$ and 10 lattices.

S. Borsanyi *et al.* "The QCD equation of state with dynamical quarks,"
JHEP **1011**, 077 (2010)



Hagedorn resonance gas: hadrons with finite widths

Modified in medium properties


$$A(M, m) \sim \frac{\Gamma \cdot m}{(M^2 - m^2)^2 + \Gamma^2 \cdot m^2} ,$$

$$\Gamma(T) = C_\Gamma \left(\frac{m}{T_H} \right)^{N_m} \left(\frac{T}{T_H} \right)^{N_T} \exp \left(\frac{m}{T_H} \right)$$

The energy density

$$\begin{aligned} \varepsilon(T, \mu_B, \mu_S) &= \sum_{i: m_i < m_0} g_i \varepsilon_i(T, \mu_i; m_i) \\ &+ \sum_{i: m_i \geq m_0} g_i \int_{m_0^2}^{\infty} d(M^2) A(M, m_i) \varepsilon_i(T, \mu_i; M), \end{aligned}$$

Hagedorn resonance gas: hadrons with finite widths



$$P(T) = T \int_0^T dT' \frac{\varepsilon(T')}{T'^2}.$$

N_m in the range from $N_m = 2.5$ (dashed line) to $N_m = 3.0$ (solid line).

$$C_\Gamma = 10^{-4}$$

$$N_T = 6.5$$

$$T_H = 165 \text{ MeV}$$

$$\Gamma(T) = C_\Gamma \left(\frac{m}{T_H} \right)^{N_m} \left(\frac{T}{T_H} \right)^{N_T} \exp\left(\frac{m}{T_H} \right)$$

D. Blaschke & K.A. Bugaev, Fizika B **13**, 491 (2004); PPNP **53**, 197 (2004)



Mott-Hagedorn resonance gas

State-dependent hadron resonance width


$$A_i(M, m_i) \sim \frac{\Gamma_i \cdot m_i}{(M^2 - m_i^2)^2 + \Gamma_i^2 \cdot m_i^2},$$

$$\Gamma_i(T) = \tau_{\text{coll},i}^{-1}(T) = \sum_j \lambda \langle r_i^2 \rangle_T \langle r_j^2 \rangle_T n_j(T)$$

D. B. Blaschke, J. Berdermann, J. Cleymans, K. Redlich: [arXiv:1102.2908]

For mesons


$$r_\pi^2(T) = \frac{3M_\pi^2}{4\pi^2 m_q} |\langle \bar{q}q \rangle_T|^{-1}; \quad \langle \bar{q}q \rangle_T \sim [1 - \tanh(0.002T - 1)]$$



For baryons: $r_N^2(T, \mu) = r_0^2 + r_\pi^2(T, \mu)$ with $r_0 = 0.45 \text{ fm}$.



Mott-Hagedorn resonance gas



Mott-Hagedorn resonance gas:

Pressure and energy density for three values of the mass threshold

$m_0 = 1.0$ GeV (solid lines)

$m_0 = 0.98$ GeV (dashed lines) and

$m_0 = 0$ (dash-dotted lines)

$$\varepsilon(T) = \sum_{i: m_i < m_0} g_i \varepsilon_i(T; m_i) + \sum_{i: m_i \geq m_0} g_i \int_{m_0^2}^{\infty} d(M^2) A(M, m_i) \varepsilon_i(T; M),$$

Quarks and gluons are missing!



Quarks and gluons in the PNJL model

Systematic expansion of the pressure as the thermodynamical potential in the grand canonical ensemble for a chiral quark model of the PNJL type beyond its mean field description $P_{PNJL, MF}(T)$ by including perturbative with α_S corrections

$$P(T) = P_{MHRG}(T) + P_{PNJL, MF}(T) + P_2(T) .$$

$P_{MHRG}(T)$ stands for the pressure of the MHRG model, accounting for the dissociation of hadrons in hot dense matter.

Quark and gluon contributions

$$P_2(T) = P_2^{\text{quark}}(T) + P_2^{\text{gluon}}(T)$$



Quark and gluon contributions

$P_2^{\text{quark}}(T)$



$P_2^{\text{gluon}}(T)$



Total perturbative QCD correction

$$P_2 = -\frac{8}{\pi} \alpha_s T^4 (I_\Lambda^+ +$$

$$\frac{3}{\pi^2} ((I_\Lambda^+)^2 + (I_\Lambda^-)^2))$$

$$\xrightarrow{\Lambda/T \rightarrow 0} -\frac{3\pi}{2} \alpha_s T^4$$

where

$$I_\Lambda^\pm = \int_{\Lambda/T}^{\infty} \frac{dx x}{e^x \pm 1}$$


Energy corrections

$$\varepsilon_2(T) = T \frac{dP_2(T)}{dT} - P_2(T)$$



Quarks, gluons and hadron resonances

$$P_{\text{MHRG}}(T) = \sum_i \delta_i d_i \int \frac{d^3 p}{(2\pi)^3} \int dM A_i(M, m_i) T \ln \left\{ 1 + \delta_i e^{-[\sqrt{p^2 + M^2} - \mu_i]/T} \right\},$$




- Quark-gluon plasma contributions are described within the improved PNJL model with α_s corrections.
- Heavy hadrons are described within the resonance gas with finite width exhibiting a Mott effect at the coincident chiral and deconfinement transitions.



Further progress

Taking into account hadrons scattering in the dense medium:

The generalized Beth-Uhlenbeck approach and construction of the scattering phase shift taking into account Levinson theorem (A. Wergieluk, D. Blaschke & Co.)



Conclusions

- An effective model description of QCD thermodynamics at finite temperatures which properly accounts for the fact that in the QCD transition region it is dominated by a tower of hadronic resonances.
- A generalization of the Hagedorn resonance gas thermodynamics which includes the finite lifetime of hadronic resonances in a hot and dense medium

