

Low mass dilepton production in pp, p-Pb and Pb-Pb collisions measured with ALICE at the LHC

Tuesday, 5 November 2013 13:50 (20 minutes)

Low mass dilepton production, including light vector mesons ρ , ω , ϕ , provides key information on the hot and dense state of strongly interacting matter produced in high-energy heavy-ion collisions. Among them, strangeness production can be studied through the measurement of ϕ meson production, while the detailed description of the full dilepton mass spectra down to the kinematic threshold can be used to reveal in-medium modifications of hadron properties and the thermal emission arising from the medium. Measurements in pp and p-A systems, in absence of hot nuclear matter effects, can be used as a reference to test our knowledge of the processes expected to contribute to dilepton production.

Dilepton production is studied with the ALICE apparatus at the LHC both at central ($|y| < 0.9$) and forward ($2.5 < y < 4$) rapidities, respectively in the dielectron and dimuon channel. Results on low mass dilepton production are shown, for various c.m. energies, in pp, p-Pb, Pb-p and Pb-Pb collisions.

Primary author: URAS, Antonio (Universite Claude Bernard-Lyon I)

Presenter: URAS, Antonio (Universite Claude Bernard-Lyon I)

Session Classification: Hard and Thermal Electroweak Probes

Track Classification: Hard and Thermal Electroweak Probes