

# Measuring the gluon distribution in nuclei at an Electron-Ion Collider

Matthew A. C. Lamont  
BNL

# Lots of work recently on the physics of $e+A$ collisions

The EIC Science case:  
a report on the joint  
BNL/INT/JLab program

Gluons and the quark sea at high energies:  
distributions, polarization, tomography

Institute for Nuclear Theory • University of Washington, USA  
September 13 to November 19, 2010



Editors:

D. Boer  
Rijksuniversiteit Groningen, The Netherlands

M. Diehl  
Deutsches Elektronen-Synchrotron DESY, Germany

R. Milner  
Massachusetts Institute of Technology, USA

R. Venugopalan  
Brookhaven National Laboratory, USA

W. Vogelsang  
Universität Tübingen, Germany

[arXiv:1108.1713](https://arxiv.org/abs/1108.1713)

Hard Probes 2013: [macl@bnl.gov](mailto:macl@bnl.gov)

# Lots of work recently on the physics of $e+A$ collisions

The EIC Science case:  
a report on the joint  
BNL/INT/JLab program

**Glueons and the quark sea at high energies:  
distributions, polarization, tomography**

Institute for Nuclear Theory • University of Washington, USA  
September 13 to November 19, 2010



Editors:  
D. Boer  
Rijksuniversiteit Groningen, The Netherlands  
M. Diehl  
Deutsches Elektronen-Synchrotron DESY, Germany  
R. Milner  
Massachusetts Institute of Technology, USA

R. Venugopalan  
Brookhaven National Laboratory, USA  
W. Vogelsang  
Universität Tübingen, Germany

[arXiv:1108.1713](https://arxiv.org/abs/1108.1713)

**Electron Ion Collider:  
The Next QCD Frontier**


Understanding the glue  
that binds us all

[arXiv:1212.1701](https://arxiv.org/abs/1212.1701)

Hard Probes 2013: [macl@bnl.gov](mailto:macl@bnl.gov)




# Why $e+A$ collisions and not $p+A$ ?

- $e+A$  and  $p+A$  provide excellent information on properties of gluons in the nuclear wave functions
- Both are **complementary** and offer the opportunity to perform stringent checks of **factorization/universality**
- Issues:
  - ➔  $p+A$  combines initial and final state effects
  - ➔ multiple colour interactions in  $p+A$
  - ➔  $p+A$  lacks the direct access to  $x$ ,  $Q^2$



# Why e+A collisions and not p+A?

- e+A and p+A provide excellent information on properties of gluons in the nuclear wave functions
- Both are **complementary** and offer the opportunity to perform stringent checks of **factorization/universality**
- Issues:
  - ➔ p+A combines initial and final state effects
  - ➔ multiple colour interactions in p+A
  - ➔ p+A lacks the direct access to  $x$ ,  $Q^2$



$p_T - x$  correlation in p+p

# Why e+A collisions and not p+A?

- e+A and p+A provide excellent information on properties functions



- Both are opportunities for factorization

- Issues:

- p+A collisions
- multiple interactions
- p+A lacks

*Scattering of protons on protons is like colliding Swiss watches to find out how they are built.*

R. Feynman




$p_T \propto$  concentration in p+p

# What is eRHIC?


## Electron accelerator

(to be built)

Unpolarized and polarized leptons  
5-20 (30) GeV



70% e<sup>-</sup> beam polarization goal  
polarized positrons?




## RHIC

Existing = \$2B


Polarized protons  
50-250 GeV

Light ions (d, Si, Cu)  
Heavy ions (Au, U)  
50-100 GeV/u


Polarized light ions He<sup>3</sup>  
166 GeV/u




Center mass energy range:  $\sqrt{s}=30-200$  GeV;  $L \sim 100-1000 \times \text{Hera}$   
longitudinal and transverse polarization for p/He<sup>3</sup> possible



# From RHIC to eRHIC





# From RHIC to eRHIC



# Staging of eRHIC: $E_e$ : 5 to 30 GeV

All energies scale proportionally by adding SRF cavities to the injector



$E/E_0$
0.0200
0.1017
0.1833
0.2650
0.3467
0.4283
0.5100
0.5917
0.6733
0.7550
0.8367
0.9183
1.0000


All magnets would be installed from the day one and we would be cranking power supplies up as energy is increasing

# What did we learn from e+p collisions at HERA?

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$


quark+anti-quark  
momentum distributions

gluon momentum  
distribution



# What did we learn from e+p collisions at HERA?


$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$




Scaling violation:  $dF_2/d\ln Q^2$  and linear DGLAP Evolution  $\Rightarrow G(x, Q^2)$

# What did we learn from e+p collisions at HERA?

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$




Scaling violation:  $dF_2/d\ln Q^2$  and linear DGLAP Evolution  $\Rightarrow G(x, Q^2)$




# What did we learn from e+p collisions at HERA?

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$




Scaling violation:  $dF_2/d\ln Q^2$  and linear DGLAP Evolution  $\Rightarrow G(x, Q^2)$




# What did we learn from e+p collisions at HERA?



$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$




Scaling violation:  $dF_2/d\ln Q^2$  and linear DGLAP Evolution  $\Rightarrow G(x, Q^2)$





# The structure of matter at small-x




- Gluons dominate the PDFs at small- to intermediate- $x$  ( $x < 0.1$ )
  - ➔ Rapid rise in gluons described naturally by linear pQCD evolution equations





# The structure of matter at small-x



- Gluons dominate the PDFs at small- to intermediate- $x$  ( $x < 0.1$ )
  - ➔ Rapid rise in gluons described naturally by linear pQCD evolution equations
  - ➔ This rise cannot increase forever - limits on the cross-section
    - non-linear pQCD evolution equations provide a natural way to tame this growth and lead to a saturation of gluons, characterised by the saturation scale  $Q_s^2(x)$



# The structure of matter at small-x





- Gluons dominate the PDFs at small- to intermediate- $x$  ( $x < 0.1$ )
  - ➔ Rapid rise in gluons described naturally by linear pQCD evolution equations
  - ➔ This rise cannot increase forever - limits on the cross-section
    - non-linear pQCD evolution equations provide a natural way to tame this growth

**however - saturation in the gluon density is not observed in the gluon distribution at HERA -> too high an  $x$**


**How can this be observed at eRHIC?**

# Nuclear “oomph” effect

Pocket formula:  $Q_s^2(x) \sim A^{1/3} \left(\frac{1}{x}\right)^\lambda \sim \left(\frac{A}{x}\right)^{1/3}$




# What do we know about the structure of nuclei?



The distribution of valence and sea quarks are relatively well known in nuclei - theories agree well

# What do we know about the structure of nuclei?



The distribution of valence and sea quarks are relatively well known in nuclei - theories agree well

Large discrepancies exist in the gluon distributions from models for mid-rapidity LHC and forward RHIC rapidities !!

# Phase-space coverage of e+A collisions for an EIC

- Existing data:

- ➔ Low energy (fixed target)

- ➔ Low statistics


- ➔ Mainly light A

- EIC coverage:

- ➔ Both “low energy” and “high energy” options extend the reach in x-Q<sup>2</sup> beyond current data

- ➔ A coverage extended up to U

- ➔ Saturation scale at moderate Q<sup>2</sup> can be investigated at the lowest x




# Phase-space coverage of e+A collisions for an EIC

- Existing data:

- ➔ Low energy (fixed target)
- ➔ Low statistics
- ➔ Mainly light A

- EIC coverage:

- ➔ Both “low energy” and “high energy” options extend the reach in x-Q<sup>2</sup> beyond current data
- ➔ A coverage extended up to U
- ➔ Saturation scale at moderate Q<sup>2</sup> can be investigated at the lowest x




# Phase-space coverage of e+A collisions for an EIC

- Existing data:

- ➔ Low energy (fixed target)
- ➔ Low statistics
- ➔ Mainly light A

- EIC coverage:

- ➔ Both “low energy” and “high energy” options extend the reach in x-Q<sup>2</sup> beyond current data
- ➔ A coverage extended up to U
- ➔ Saturation scale at moderate Q<sup>2</sup> can be investigated at the lowest x




# Phase-space coverage of e+A collisions for an EIC

- Existing data:

- ➔ Low energy (fixed target)
- ➔ Low statistics
- ➔ Mainly light A

- EIC coverage:

- ➔ Both “low energy” and “high energy” options extend the reach in x-Q<sup>2</sup> beyond current data
- ➔ A coverage extended up to U
- ➔ Saturation scale at moderate Q<sup>2</sup> can be investigated at the lowest x




# Saturation effects in the proton and nucleus

$$\frac{d^2\sigma^{eA \rightarrow eX}}{dx dQ^2} = \frac{4\pi\alpha^2}{xQ^4} \left[ \left(1 - y + \frac{y^2}{2}\right) F_2(x, Q^2) - \frac{y^2}{2} F_L(x, Q^2) \right]$$

quark+anti-quark gluon

## Measure of non-linear effects in the $F_L$ structure function

Dipole model (J. Bartels *et al.*)



- Plotting this distribution coming out of saturation inspired GBW model
- ➔ p: small effect only starting to come in at small-x and small  $Q^2$



# Saturation effects in the proton and nucleus

$$\frac{d^2\sigma^{eA\rightarrow eX}}{dx dQ^2} = \frac{4\pi\alpha^2}{xQ^4} \left[ \left(1 - y + \frac{y^2}{2}\right) F_2(x, Q^2) - \frac{y^2}{2} F_L(x, Q^2) \right]$$

↙ quark+anti-quark      ↘ gluon

Measure of non-linear effects in the  $F_L$  structure function

Dipole model (J. Bartels *et al.*)



- Plotting this distribution coming out of saturation inspired GBW model
  - ➔ p: small effect only starting to come in at small-x and small  $Q^2$
  - ➔ Au: much larger effects are visible
    - nuclear “oomph” effects well manifested in the  $F_L$  structure function



# Saturation effects in the proton and nucleus

$$\frac{d^2\sigma^{eA \rightarrow eX}}{dx dQ^2} = \frac{4\pi\alpha^2}{xQ^4} \left[ \left(1 - y + \frac{y^2}{2}\right) F_2(x, Q^2) - \frac{y^2}{2} F_L(x, Q^2) \right]$$

↙ quark+anti-quark      ↘ gluon

Measure of non-linear effects in the  $F_L$  structure function

Dipole model (J. Bartels *et al.*)



- Plotting this distribution coming out of saturation inspired GBW model
  - ➔ p: small effect only starting to come in at small-x and small  $Q^2$
  - ➔ Au: much larger effects are visible
    - nuclear “oomph” effects well manifested in the  $F_L$  structure function

# Feasibility study:

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$

## Strategies:

slope of  $y^2/Y_+$  for different  $s$  at fixed  $x$  &  $Q^2$

## e+Au:

20x50 -  $AfLdt = 2 \text{ fb}^{-1}$

20x75 -  $AfLdt = 4 \text{ fb}^{-1}$

20x100 -  $AfLdt = 4 \text{ fb}^{-1}$

running combined










~6 months total running

(50% eff)

statistical errors are swamped by the 3% systematic errors

Will be dominated by systematics, but would need a full detector simulation in order to estimate them

$Q^2 = 1.389 \text{ GeV}^2$



# Feasibility study:

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$

## Strategies:

slope of  $y^2/Y_+$  for different  $s$  at fixed  $x$  &  $Q^2$

## e+Au:

20x50 -  $AfLdt = 2 \text{ fb}^{-1}$

20x75 -  $AfLdt = 4 \text{ fb}^{-1}$

20x100 -  $AfLdt = 4 \text{ fb}^{-1}$


running combined

~6 months total running  
(50% eff)

statistical errors are  
swamped by the 3%  
systematic errors

Will be dominated by  
systematics, but would  
need a full detector  
simulation in order to  
estimate them

Fixed  $F_2$  (0.5) and  $F_L$  (0.2)



# Feasibility study:

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$

## Strategies:

slope of  $y^2/Y_+$  for different  $s$  at fixed  $x$  &  $Q^2$

## e+Au:

20x50 -  $AfLdt = 2 \text{ fb}^{-1}$

20x75 -  $AfLdt = 4 \text{ fb}^{-1}$

20x100 -  $AfLdt = 4 \text{ fb}^{-1}$


running combined

~6 months total running  
(50% eff)

statistical errors are  
swamped by the 3%  
systematic errors

Will be dominated by  
systematics, but would  
need a full detector  
simulation in order to  
estimate them

Fixed  $F_2$  (0.5) and  $F_L$  (0.2)



# Feasibility study:

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$

## Strategies:

slope of  $y^2/Y_+$  for different  $s$  at fixed  $x$  &  $Q^2$

## e+Au:

20x50 -  $AfLdt = 2 \text{ fb}^{-1}$

20x75 -  $AfLdt = 4 \text{ fb}^{-1}$

20x100 -  $AfLdt = 4 \text{ fb}^{-1}$


running combined

~6 months total running  
(50% eff)

statistical errors are  
swamped by the 3%  
systematic errors

Will be dominated by  
systematics, but would  
need a full detector  
simulation in order to  
estimate them

Fixed  $F_2$  (0.5) and  $F_L$  (0.2)



# Feasibility study:

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$

## Strategies:

slope of  $y^2/Y_+$  for different  $s$  at fixed  $x$  &  $Q^2$

## e+Au:

20x50 -  $AfLdt = 2 \text{ fb}^{-1}$

20x75 -  $AfLdt = 4 \text{ fb}^{-1}$

20x100 -  $AfLdt = 4 \text{ fb}^{-1}$


running combined

~6 months total running  
(50% eff)

statistical errors are  
swamped by the 3%  
systematic errors

Will be dominated by  
systematics, but would  
need a full detector  
simulation in order to  
estimate them

Fixed  $F_2$  (0.5) and  $F_L$  (0.2)



# Feasibility study:

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$

## Strategies:

slope of  $y^2/Y_+$  for different  $s$  at fixed  $x$  &  $Q^2$

## e+Au: 1st stage

5x50 -  $AfLdt = 2 \text{ fb}^{-1}$

5x75 -  $AfLdt = 4 \text{ fb}^{-1}$



5x100 -  $AfLdt = 4 \text{ fb}^{-1}$

running combined

~6 months total running  
(50% eff)

statistical errors are swamped by the 1% systematic errors

Will be dominated by systematics, but would need a full detector simulation in order to estimate them




# Work in progress... (H. Paukkunen)

- Take the generated Pseudo-data and include it in a global fit

➔ Only 20x100 and 5x100 included in these plots

- More data (e.g. charm) will constrain this further




# Work in progress... (H. Paukkunen)

- Take the generated Pseudo-data and include it in a global fit

➔ Only 20x100 and 5x100 included in these plots

- More data (e.g. charm) will constrain this further




# Work in progress... (H. Paukkunen)

- Take the generated Pseudo-data and include it in a global fit

➔ Only 20x100 and 5x100 included in these plots

- More data (e.g. charm) will constrain this further



# Summary and Conclusions

- The **e+A physics programme** at an **EIC** will give us an unprecedented opportunity to study gluons in nuclei
  - ➔ **Low-x:** Measure the properties of gluons where saturation is the dominant governing phenomena
  - ➔ **Higher-x:** Understand how fast partons interact as they traverse nuclear matter and provide new insight into hadronization

# Summary and Conclusions

- The **e+A physics programme** at an **EIC** will give us an unprecedented opportunity to study gluons in nuclei
  - ➔ **Low-x:** Measure the properties of gluons where saturation is the dominant governing phenomena
  - ➔ **Higher-x:** Understand how fast partons interact as they traverse nuclear matter and provide new insight into hadronization
- Understanding gluons' role in nuclei is crucial to understanding RHIC and LHC results
  - ➔ Much information obtained from measuring the  $F_2$  and  $F_L$  structure functions

# Summary and Conclusions

- The **e+A physics programme** at an **EIC** will give us an unprecedented opportunity to study gluons in nuclei
  - ➔ **Low-x**: Measure the properties of gluons where saturation is the dominant governing phenomena
  - ➔ **Higher-x**: Understand how fast partons interact as they traverse nuclear matter and provide new insight into hadronization
- Understanding gluons' role in nuclei is crucial to understanding RHIC and LHC results
  - ➔ Much information obtained from measuring the  $F_2$  and  $F_L$  structure functions
- Other important observables not discussed due to time limitations:
  - ➔ **Diffraction**: The diffractive cross-section itself gives information on the level of saturation
  - ➔ **di-Hadron Correlations**: Analogue measurement to p/d+A, but less uncertainties on the measurement

# Summary and Conclusions

- The **e+A physics programme** at an **EIC** will give us an unprecedented opportunity to study gluons in nuclei
  - ➔ **Low-x**: Measure the properties of gluons where saturation is the dominant governing phenomena
  - ➔ **Higher-x**: Understand how fast partons interact as they traverse nuclear matter and provide new insight into hadronization
- Understanding gluons' role in nuclei is crucial to understanding RHIC and LHC results
  - ➔ Much information obtained from measuring the  $F_2$  and  $F_L$  structure functions
- Other important observables not discussed due to time limitations:
  - ➔ **Diffraction**: The diffractive cross-section itself gives information on the level of saturation
  - ➔ **di-Hadron Correlations**: Analogue measurement to p/d+A, but less uncertainties on the measurement
- The INT programme in the Autumn of 2010 allowed us to formulate the observables in terms of golden and silver measurements
  - ➔ A detailed write-up of the whole programme is on the **ArXiv: 1108.1713**
  - ➔ An EIC White Paper (not just e+A), expounding on the INT programme has been released to the community **ArXiv: 1212.1701**

# Summary and Conclusions


- The **e+A physics programme** at an **EIC** will give us an unprecedented opportunity to study gluons in nuclei
  - ➔ **Low-x**: Measure the properties of gluons where saturation is the dominant governing phenomena
  - ➔ **Higher-x**: Understand how fast partons interact as they traverse nuclear matter and provide new insight into hadronization

**entire science programme is uniquely tied to a  
future high-energy electron-ion collider  
never been measured before & never without**

- ➔ **di-Hadron Correlations**: Analogue measurement to p/d+A, but less uncertainties on the measurement
- The INT programme in the Autumn of 2010 allowed us to formulate the observables in terms of golden and silver measurements
  - ➔ A detailed write-up of the whole programme is on the **ArXiv: 1108.1713**
  - ➔ An EIC White Paper (not just e+A), expounding on the INT programme has been released to the community **ArXiv: 1212.1701**

**BACKUP**

# Exclusive processes in e+A - diffraction



- $\beta$  is the momentum fraction of the struck parton w.r.t. the Pomeron
- $x_{IP} = x/\beta$ : momentum fraction of the exchanged object (Pomeron) w.r.t. the hadron

$$\beta = \frac{x}{x_{IP}} = \frac{Q^2}{Q^2 + M_X^2 - t}$$





## • Diffraction in e+p:

- ➔ HERA: 15% of all events are diffractive

## • Diffraction in e+A:


- ➔ Predictions:  $\sigma_{\text{diff}}/\sigma_{\text{tot}}$  in e+A ~25-40%
- ➔ Coherent diffraction (nuclei intact)
- ➔ Incoherent diffraction: breakup into nucleons (nucleons intact)

# Day 1: Diffractive Cross-sections




- **Ratio of diffractive-to-total cross-section** drastically different between saturation (Marquet) and non-saturation (Frankfurt, Guzey, Strikman) models
- Expected experimental error bars (simulated for  $10 \text{ fb}^{-1}$  of data for a **low-energy eRHIC**) can distinguish between the two scenarios

# Exclusive vector meson production




$$d\sigma \propto g(x)^2$$

Sartre: Toll, Ullrich (2012)




- Exclusive vector meson production is most sensitive to the gluon distribution
  - ➔ colour-neutral exchange of gluons
- J/ψ shows some difference between saturation and no-saturation

# Exclusive vector meson production





$$d\sigma \propto g(x)^2$$

Sartre: Toll, Ullrich (2012)



- Exclusive vector meson production is most sensitive to the gluon distribution
  - ➔ colour-neutral exchange of gluons
- $J/\psi$  shows some difference between saturation and no-saturation
- $\phi$  shows a much larger difference
  - ➔ wave function for  $\phi$  is larger and hence more sensitive to saturation effects

# Exclusive Vector Meson Production in e+A





- Low- $t$ : coherent diffraction dominates - gluon density
  - High- $t$ : incoherent diffraction dominates - gluon correlations
- ➔ Need good breakup detection efficiency to discriminate between the two scenarios
- unlike protons, forward spectrometer won't work for heavy ions
  - measure emitted neutrons in a ZDC
  - rapidity gap with absence of break-up fragments sufficient to identify coherent events

# Finding the source...

- Take the  $d\sigma/dt$  distribution and perform a Fourier

$$F(b) \sim \frac{1}{2\pi} \int_0^{\infty} d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$  (for small  $x$ )





# Finding the source...

- Take the  $d\sigma/dt$  distribution and perform a Fourier

$$F(b) \sim \frac{1}{2\pi} \int_0^{\infty} d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$  (for small  $x$ )





# Finding the source...

- Take the  $d\sigma/dt$  distribution and perform a Fourier

$$F(b) \sim \frac{1}{2\pi} \int_0^{\infty} d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$  (for small  $x$ )





# Finding the source...

- Take the  $d\sigma/dt$  distribution and perform a Fourier

$$F(b) \sim \frac{1}{2\pi} \int_0^{\infty} d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$  (for small x)





# Finding the source...

- Take the  $d\sigma/dt$  distribution and perform a Fourier

$$F(b) \sim \frac{1}{2\pi} \int_0^{\infty} d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$  (for small  $x$ )





# Finding the source...

- Take the  $d\sigma/dt$  distribution and perform a Fourier

$$F(b) \sim \frac{1}{2\pi} \int_0^{\infty} d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$  (for small  $x$ )





# Finding the source...

- Take the  $d\sigma/dt$  distribution and perform a Fourier

$$F(b) \sim \frac{1}{2\pi} \int_0^{\infty} d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$  (for small  $x$ )





# Finding the source...

- Take the  $d\sigma/dt$  distribution and perform a Fourier





$$F(b) \sim \frac{1}{2\pi} \int_0^{\infty} d\Delta \Delta J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}}$$

$t = \Delta^2/(1-x) \approx \Delta^2$  (for small  $x$ )




# Finding the source...

- $J/\psi$  shows little difference for both saturated and non-saturated modes.



# di-hadron correlations in d+A


comparisons between  $d+Au \rightarrow h_1 h_2 X$  (or  $p+Au \rightarrow h_1 h_2 X$ ) and  $p+p \rightarrow h_1 h_2 X$



- At  $y=0$ , suppression of away-side jet is observed in A+A collisions
- No suppression in p+p or d+A


$\Rightarrow x \sim 10^{-2}$

$$x_A = \frac{k_1 e^{-y_1} + k_2 e^{-y_2}}{\sqrt{s}} \ll 1$$



# di-hadron correlations in d+A


comparisons between d+Au  $\rightarrow h_1 h_2 X$  (or p +Au  $\rightarrow h_1 h_2 X$ ) and p+p  $\rightarrow h_1 h_2 X$





- At  $y=0$ , suppression of away-side jet is observed in A+A collisions
- No suppression in p+p or d+A

$\rightarrow x \sim 10^{-2}$

$$x_A = \frac{k_1 e^{-y_1} + k_2 e^{-y_2}}{\sqrt{s}} \ll 1$$



- However, at forward rapidities ( $y \sim 3.1$ ), an away-side suppression is observed in d+Au





- Away-side peak



[bnl.gov](http://bnl.gov)

# di-hadron correlations in d+A

comparisons b  
+Au  $\rightarrow$   $h_1 h_2$



$p+p \rightarrow \pi^0 \pi^0 + X$   
 $p_{T,L} > 2 \text{ GeV}/c, 1 < \eta_L <$



of away-  
in A+A

+p or d+A

$$\frac{k_2 e^{-y_2}}{\sqrt{s}} \ll 1$$


at  
pseudorapidities ( $y$ )

away-  
emission is  
in d+Au

peak

# di-hadron correlations in e+A

Never been measured - we expect to see the same effect in e+A as in d+A




Dominguez, Xiao and Yuan (2012)

- At small-x, multi-gluon distributions are as important as single-gluon distributions and they contribute to di-hadron correlations
  - ➔ The non-linear evolution of multi-gluon distributions is different from that of single-gluon distributions and it is **equally important** that we understand it
- The d+Au RHIC data is therefore subject to many uncertainties
  - ➔ these correlations in e+A can help to constrain them better

# di-hadron correlations in e+A

Never been measured - we expect to see the same effect in e+A as in d+A



Dominguez, Xiao and Yuan (2012)

- At small-x, multi-gluon distributions are as important as single-gluon distributions and they contribute to di-hadron correlations

➔ The non-linear evolution of multi-gluon distributions is different from that of single-gluon distributions and it is **equally important** that we understand it



- The d+Au RHIC data is therefore subject to many uncertainties

➔ these correlations in e+A can help to constrain them better

# di-hadron Correlations - relative yields


- PHENIX measured  $J_{dAu}$  - relative yield of di-hadrons produced in d+Au compared to p+p collisions

➔ Suppression in central events compared to peripheral as a function of  $x_A^{\text{frag}}$




A. Adare et al., Phys. Rev. Lett. 107, 172301 (2011)

# Jets and hadronization



- $t_p$  - production time of propagating quark
- $t_f^h$  - hadron formation time


# Jets and hadronization



What happens if we add a nuclear medium?

- $t_p$  - production time of propagating quark
- $t_f^h$  - hadron formation time


# Jets and hadronization



What happens if we add a nuclear medium?

- $t_p$  - production time of propagating quark
- $t_f^h$  - hadron formation time

# Jets and hadronization




What happens if we add a nuclear medium?

- $t_p$  - production time of propagating quark
- $t_f^h$  - hadron formation time

## Observables:

Broadening:  $\Delta p_T^2 = \langle p_T^2 \rangle_A - \langle p_T^2 \rangle_p$ : direct link to saturation scale

# Jets and hadronization



What happens if we add a nuclear medium?


- $t_p$  - production time of propagating quark
- $t_f^h$  - hadron formation time

## Observables:

Broadening:  $\Delta p_T^2 = \langle p_T^2 \rangle_A - \langle p_T^2 \rangle_p$  : direct link to saturation scale

Attenuation:  $R_A^h(Q^2, \nu, z_h, p_T^2)$  : ratio of hadron production in A to D, modifications of nPDFs cancel out

# Jets and hadronization



- $t_p$  - production time of propagating quark
- $t^{h_f}$  - hadron formation time

## Observables:

Broadening:  $\Delta p_T^2 = \langle p_T^2 \rangle_A - \langle p_T^2 \rangle_p$  : direct link to saturation scale


Attenuation:  $R_A^h(Q^2, \nu, z_h, p_T^2)$  : ratio of hadron production in A to D,  
modifications of nPDFs cancel out

# How can the EIC contribute?

HERMES:

$$E_e = 27 \text{ GeV} \rightarrow \sqrt{s} = 7.2 \text{ GeV}$$

$$E_h = 2\text{-}15 \text{ GeV}$$



$v$  = virtual photon energy

$$z_h = E_h/v$$



Hard Probes 2013: [macl@bnl.gov](mailto:macl@bnl.gov)

# How can the EIC contribute?

HERMES:

$$E_e = 27 \text{ GeV} \rightarrow \sqrt{s} = 7.2 \text{ GeV}$$

$$E_h = 2\text{-}15 \text{ GeV}$$



$\nu$  = virtual photon energy

$$Z_h = E_h/\nu$$


Hard Probes 2013: [macl@bnl.gov](mailto:macl@bnl.gov)

# How can the EIC contribute?

HERMES:

$$E_e = 27 \text{ GeV} \rightarrow \sqrt{s} = 7.2 \text{ GeV}$$

$$E_h = 2\text{-}15 \text{ GeV}$$





$\nu$  = virtual photon energy

$$Z_h = E_h/\nu$$

EIC:

light hadrons:




large  $\nu$  range  $\rightarrow$  boost

# How can the EIC contribute?

HERMES:

$$E_e = 27 \text{ GeV} \rightarrow \sqrt{s} = 7.2 \text{ GeV}$$

$$E_h = 2\text{-}15 \text{ GeV}$$




$\nu$  = virtual photon energy

$$Z_h = E_h/\nu$$

EIC:

charm hadrons:



large  $\nu$  range  $\rightarrow$  boost

Hard Probes 201 hadronization in and out of nucleus 29


# How can the EIC contribute?

HERMES:

$$E_e = 27 \text{ GeV} \rightarrow \sqrt{s} = 7.2 \text{ GeV}$$

$$E_h = 2\text{-}15 \text{ GeV}$$

EIC:




$\nu$  = virtual photon energy

$$Z_h = E_h/\nu$$

large  $\nu$  range  $\rightarrow$  boost

# Jets at an EIC



- E665 at FNAL have measured jets in  $\mu+A$  at  $\sqrt{s} \sim 30$  GeV
  - ➔ Feasible to start a jet programme in phase 1
  - ➔ caveat that collider kinematics are different to fixed target



# Jets at an EIC


- E665 at FNAL have measured jets in  $\mu+A$  at  $\sqrt{s} \sim 30$  GeV
  - ➔ Feasible to start a jet programme in phase 1
  - ➔ caveat that collider kinematics are different to fixed target

1+1 jets, dominated by q processes → allow study of parton propagation through cold nuclear matter




# Jets at an EIC

- E665 at FNAL have measured jets in  $\mu+A$  at  $\sqrt{s} \sim 30$  GeV
  - ➔ Feasible to start a jet programme in phase 1
  - ➔ caveat that collider kinematics are different to fixed target





1+1 jets, dominated by q processes → allow study of parton propagation through cold nuclear matter



$$\frac{d^2\sigma_{2+1}}{dx dQ^2} = A_q(x, Q^2)q^A(x, Q^2) + A_g(x, Q^2)g_A(x, Q^2)$$


2+1 jets → sensitive to nuclear gluons



By measuring 1+1 jets, can extract information on gluons



# Jets at an EIC

1+1 jets, dominated by q processes → allow study of parton propagation through cold nuclear matter





$$\frac{d^2\sigma_{2+1}}{dx dQ^2} = A_q(x, Q^2)q^A(x, Q^2) + A_g(x, Q^2)g_A(x, Q^2)$$



2+1 jets → sensitive to nuclear gluons



By measuring 1+1 jets, can extract information on gluons



# What do we know about the structure of nuclei?



- e+p data covers large part of phase space

➔ low  $x$  and large  $Q^2$

- e+A data only a small fraction of this (e+A was a fixed target programme at HERA)

➔ high-medium  $x$  and low  $Q^2$

# $F_L$ at an EIC vs $F_L$ at LHeC





Fig 6.18 of LHEC CDR


- $R_{F_2}$  has good coverage with  $x$  and small uncertainties

# $F_L$ at an EIC vs $F_L$ at LHeC




- $R_{F_2}$  has good coverage with  $x$  and small uncertainties
- $R_{F_L}$  concentrated at low- $x$  but large uncertainties

# $F_L$ at an EIC vs $F_L$ at LHeC



- $R_{F_2}$  has good coverage with  $x$  and small uncertainties
- $R_{F_L}$  concentrated at low- $x$  but large uncertainties

# $F_L$ at an EIC vs $F_L$ at LHeC



- $R_{F_2}$  has good coverage with  $x$  and small uncertainties
  - $R_{F_L}$  concentrated at low- $x$  but large uncertainties
- ➔ EIC data will provide constraints at higher  $x$  with smaller error bars

# Feasibility

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$

## Strategies:

slope of  $y^2/Y_+$  for different  $s$  at fixed  $x$  &  $Q^2$

## e+Au:

20x50 -  $AfLdt = 2 \text{ fb}^{-1}$

20x75 -  $AfLdt = 4 \text{ fb}^{-1}$

20x100 -  $AfLdt = 4 \text{ fb}^{-1}$










running combined

~6 months total running  
(50% eff)

statistical errors are swamped by the 3% systematic errors

Will be dominated by systematics, but would need a full detector simulation in order to estimate them

$Q^2 = 1.389 \text{ GeV}^2$



# Feasibility

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$

## Strategies:

slope of  $y^2/Y_+$  for different  $s$  at fixed  $x$  &  $Q^2$

## e+Au:

20x50 -  $AfLdt = 2 \text{ fb}^{-1}$

20x75 -  $AfLdt = 4 \text{ fb}^{-1}$

20x100 -  $AfLdt = 4 \text{ fb}^{-1}$

running combined


~6 months total running

(50% eff)

statistical errors are swamped by the 3% systematic errors

Will be dominated by systematics, but would need a full detector simulation in order to estimate them

$Q^2 = 1.389 \text{ GeV}^2$



# Feasibility

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$

## Strategies:

slope of  $y^2/Y_+$  for different  $s$  at fixed  $x$  &  $Q^2$

## e+Au:

20x50 -  $AfLdt = 2 \text{ fb}^{-1}$

20x75 -  $AfLdt = 4 \text{ fb}^{-1}$

20x100 -  $AfLdt = 4 \text{ fb}^{-1}$

running combined


~6 months total running

(50% eff)

statistical errors are swamped by the 3% systematic errors

Will be dominated by systematics, but would need a full detector simulation in order to estimate them

$Q^2 = 1.389 \text{ GeV}^2$



# Feasibility

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$

## Strategies:

slope of  $y^2/Y_+$  for different  $s$  at fixed  $x$  &  $Q^2$

## e+Au:

20x50 -  $AfLdt = 2 \text{ fb}^{-1}$

20x75 -  $AfLdt = 4 \text{ fb}^{-1}$


20x100 -  $AfLdt = 4 \text{ fb}^{-1}$

running combined

~6 months total running  
(50% eff)

statistical errors are  
swamped by the 3%  
systematic errors

Will be dominated by  
systematics, but would  
need a full detector  
simulation in order to  
estimate them



# Feasibility

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$

## Strategies:

slope of  $y^2/Y_+$  for different  
s at fixed  $x$  &  $Q^2$

## e+Au:

20x50 -  $AfLdt = 2 \text{ fb}^{-1}$

20x75 -  $AfLdt = 4 \text{ fb}^{-1}$


20x100 -  $AfLdt = 4 \text{ fb}^{-1}$

running combined

~6 months total running  
(50% eff)

statistical errors are  
swamped by the 3%  
systematic errors

Will be dominated by  
systematics, but would  
need a full detector  
simulation in order to  
estimate them



# Feasibility

$$\sigma_r(x, Q^2) = F_2^A(x, Q^2) - \frac{y^2}{Y_+} F_L^A(x, Q^2)$$

## Strategies:

slope of  $y^2/Y_+$  for different  $s$  at fixed  $x$  &  $Q^2$

## e+Au:

20x50 -  $AfLdt = 2 \text{ fb}^{-1}$

20x75 -  $AfLdt = 4 \text{ fb}^{-1}$


20x100 -  $AfLdt = 4 \text{ fb}^{-1}$

running combined


~6 months total running  
(50% eff)

statistical errors are  
swamped by the 3%  
systematic errors

Will be dominated by  
systematics, but would  
need a full detector  
simulation in order to  
estimate them




# Electron beam evolution in eRHIC's ERL




**ERL:**  
energy recovery  
linac

# Electron beam evolution in eRHIC's ERL




**ERL:**  
energy recovery  
linac

# Electron beam evolution in eRHIC's ERL




**ERL:**  
energy recovery  
linac

# Electron beam evolution in eRHIC's ERL




**ERL:**  
energy recovery  
linac

# Electron beam evolution in eRHIC's ERL



**ERL:**  
energy recovery  
linac


# Electron beam evolution in eRHIC's ERL



**ERL:**  
energy recovery  
linac




# Electron beam evolution in eRHIC's ERL



**ERL:**  
energy recovery  
linac

# Diffractive Events: Experimental Side

- How to identify
- diffractive events?



## ➔ Rapidity Gap

- requires hermetic (large acceptance) detector

## ➔ Separating coherent from incoherent diffraction

- detector and IR needs to be carefully designed to detect nuclear breakup


## ➔ Limitation at a collider

- Coherent: scattered ion cannot be measured,  $t$  not directly measurable (may be in very light ions)
- Breakup can be detected using emitted  $n$  and  $\gamma$ , some charged fragments can be measured in Roman Pots

# Large Rapidity Gap Method (LRG)

## → Identify Most Forward Going Particle (MFP)


- Works at HERA but higher  $\sqrt{s}$
- EIC smaller beam rapidities



## Hermeticity requirement:

- needs just to detector presence
- does not need momentum or PID
- simulations:  $\sqrt{s}$  not a show stopper for EIC  
(can achieve 1% contamination, 80% efficiency)

## Diffractive $\rho^0$ production at EIC: $\eta$ of MFP



# Detecting Nuclear Breakup

➔ Detecting **all** fragments  $p_{A'} = \sum p_n + \sum p_p + \sum p_d + \sum p_\alpha \dots$  not possible

➔ Focus on n emission

- Zero-Degree Calorimeter
- Requires careful design of IR

• Additional measurements:

- ▶ Fragments via Roman Pots
- ▶  $\gamma$  via EMC

Traditional modelling done in


$pA$ :

## Intra-Nuclear Cascade

- Particle production
- Remnant Nucleus ( $A, Z, E^*, \dots$ )
- ISABEL, INCL4




## De-Excitation

- Evaporation
- Fission
- Residual Nuclei
- Gemini++, SMM, ABLA (all no  $\gamma$ )



# Experimental Reality


- Here eRHIC IR layout:
- Need  $\pm X$  mrad opening through triplet for  $n$  and room for ZDC
- Big questions:
  - ➔ Excitation energy  $E^*$ ?
  - ➔ ep:  $d\sigma/M_Y \sim 1/M_Y^2$
  - ➔ eA? Assume ep and use  $E^* = M_Y - m_p$  as lower limit



is 20

# Experimental Reality

- Here eRHIC IR layout:
- Need  $\pm X$  mrad opening through triplet for  $n$  and room for ZDC
- Big questions:
  - ➔ Excitation energy  $E^*$ ?
  - ➔ ep:  $d\sigma/M_Y \sim 1/M_Y^2$
  - ➔ eA? Assume ep and use  $E^* = M_Y - m_p$  as lower limit



Simulations using Gemini++ & SMM show **it works**:

- For  $E^*_{tot} \geq 10$  MeV and 2.5 mrad  $n$  acceptance we have rejection power of at least  $10^5$ .
- Separating incoherent from coherent diffractive events is possible at a collider with  $n$ -detection via ZDCs alone