



Contribution ID: 39

Type: Oral

Statistical Properties of highly-deformed Samarium isotopes

The rare-earth isotopic chain of Samarium provides an excellent opportunity to systematically investigate the evolution of nuclear structure effects from the near spherical ($\beta_2=0.09$) ^{144}Sm isotope to the highly-deformed system ($\beta_2=0.34$) ^{154}Sm . As the nuclear shape changes, statistical properties such as the nuclear level density (NLD) and γ -strength function (γSF) are expected to be affected. In particular resonance modes, such as the Pygmy Dipole (PDR), Scissors Resonances (SR) and the recently discovered Low-Energy Enhancement (LEE) in rare-earth region may reveal interesting features when their evolution is investigated across several nuclei in an isotopic chain. Most reliable knowledge can be obtained when results from several different experiments are compared. An experiment was performed in September 2016 at Oslo Cyclotron Laboratory (OCL) where the NaI(Tl) γ -ray array, silicon particle telescopes and 6 high-efficiency LaBr₃:Ce detectors were utilized to measure particle- γ coincidence events from which the NLDs and γSFs will be extracted below the neutron separation energy threshold, S_n , using the Oslo Method (A. Schiller et al. 2000). The deuteron beam with 13 and 15 MeV energies was used to populate excited states in $^{154,155}\text{Sm}$ through the inelastic scattering ($d,d'\gamma$) and the transfer reaction (d,p). Based on the results from these measurements, the extracted NLDs and γSFs will be used to investigate the evolution of nuclear structure effects in $^{154,155}\text{Sm}$ and provide complementary information to the $^{154}\text{Sm}(p,p')^{154}\text{Sm}$ and $^{154}\text{Sm}(\alpha,\alpha')^{154}\text{Sm}$ data on resonance features that lie on the low-energy tail of the GDR. In addition, the results will further provide a near-complete picture on the evolution of the PDR, SR and/or the LEE as the isotopic chain transitions from near spherical to very deformed systems. In this talk I will present preliminary results of this investigation of statistical properties for $^{154,155}\text{Sm}$ in comparison to the previous and recent measurements of $^{148,149}\text{Sm}$ and $^{151,153}\text{Sm}$ isotopes, respectively, and ongoing measurements of $^{152,153}\text{Sm}$ at OCL.

Primary authors: Mr MALATJI, Kgashane (University of Stellenbosch); Dr WIEDEKING, Mathis (iThemba LABS); Prof. SIEM, Sunniva (University of Oslo); Mr AY, Kursad (Eskisehir Osmangazi University); Ms BECKMANN, Kristine (University of Oslo); Dr BELLO GARROTE, Frank (University of Oslo); Dr CRESPO CAMPO, Lucia (University of Oslo); Prof. ANDREAS, GÖRGEN (University of Oslo); Prof. GUTTORMSEN, Magne (University of Oslo); Dr HAGEN, Trine (University of Oslo); Mr VETLE, INGEBERG (University of Oslo); Dr JONES, Pete (iThemba LABS); Dr KHESWA, Bonginkosi (University of Johannesburg); Dr KLINTEFJORD, Malin (University of Oslo); Dr KRUGMANN, Andreas (Technische Universität Darmstadt); Dr LARSEN, Ann-Cecilie (University of Oslo); Mr MIDTBØ, Jørgen (University of Oslo); Mr OZGUR, Mustafa (Eskisehir Osmangazi University); Prof. PAPKA, Paul (University of Stellenbosch); Dr PELLEGRINI, Luna (University of the Witwatersrand); Dr RENSTRØM, Therese (University of Oslo); Dr ROSE, Sunniva (University of Oslo); Dr SAHIN, Eda (University of Oslo); Dr TVETEN, Gry (University of Oslo); Prof. VON NEUMANN-COSEL, Peter (Technische Universität Darmstadt)

Presenter: Mr MALATJI, Kgashane (University of Stellenbosch)

Track Classification: Track B