



Contribution ID: 86

Type: Oral

## Unravelling $^{12}\text{C}$ , the ISGMR and the Hoyle's ghost.

The  $^{14}\text{C}(\text{p,t})^{12}\text{C}$  reaction was employed to investigate the much-discussed excitation energy region above the Hoyle state: there is evidence that suggests that in addition to the broad  $0^+$  state at 10.3 MeV, an additional  $0^+$  resonance is required at  $\sim 8.7$  MeV. AMD calculations suggest that the Isoscalar Giant Monopole Resonance (ISGMR) may contribute to this region and suggest two distinct oscillation modes may play a role. The development of a new multi-channel R-matrix fitting programme aims to provide new insight by simultaneously analysing low-energy resonance data from multiple experiments (with different reaction channels). Moreover, a multi-channel two-level approximation is implemented which may help determine the contribution to this region from the ghost of the Hoyle state and any associated coherent interferences with other  $0^+$  states.

**Primary author:** Mr LI, Kevin Ching Wei (iThemba LABS, Stellenbosch University)

**Co-authors:** Dr SMIT, Frederick (iThemba LABS); Dr NEVELING, Retief (iThemba LABS); Prof. PAPKA, Paul (iThemba LABS, Stellenbosch University)

**Presenter:** Mr LI, Kevin Ching Wei (iThemba LABS, Stellenbosch University)

**Track Classification:** Track A