



Contribution ID: 37

Type: Oral

Investigating the influence of deformation on the low-lying electric dipole (E1) response in $^{144,154}\text{Sm}$

The past decade has seen an increase in studies dedicated to understanding the low-lying electric dipole (E1) response, commonly referred to as the Pygmy Dipole Resonance (PDR). These studies revealed that the PDR has a mixed isospin nature, and that the use of complimentary probes is needed to fully understand this response. Since majority of studies on the PDR focussed on spherical nuclei, the influence that deformation has on the PDR response is yet to be understood.

Preliminary (p,p') studies on ^{154}Sm performed at RCNP, showed potential evidence for a splitting in the PDR response similar to that of the Giant Dipole Resonance with deformation [1]. A tentative interpretation suggested that this splitting could be connected to the splitting of the resonance structure with respect to the K quantum number. In another investigation, liquid drop model calculations comparing the ratio between the transition probability of K=0 and K=1 contributions for the isovector and isoscalar components of the PDR respectively, showed that the isoscalar part varies more dramatically with deformation [2].

Therefore, studies using the $(\alpha,\alpha'\gamma)$ inelastic scattering reaction were performed on the spherical ^{144}Sm and prolate deformed ^{154}Sm at iThemba LABS. The experiments made use of the K600 magnetic spectrometer in 0° mode in co-incidence with BaGeL (Ball of High-Purity Germanium and large volume LaBr 3 :Ce detectors). In this talk, we will present the recent results of this study.

[1] A.Krugmann, PhD Thesis, (2014).

[2] M.Faccioli, J.A. Lay, A. Vitturi, M.V. Andrés, E.G. Lanza (to be published).

Primary authors: JIVAN, Harshna (University of the Witwatersrand); PELLEGRINI, L. (University of the Witwatersrand and iThemba LABS)

Co-authors: BAHINI, A. (University of the Witwatersrand); GORGEN, A. (University of Oslo); NEGRET, A. (IFIN-HH, Romania); OLACEL, A. (IFIN-HH, Romania); SAVRAN, D. (ExtreMe Matter Institute); MARIN-LAMBARRI, D.J. (iThemba LABS); SIDERAS-HADDAD, E. (University of the Witwatersrand and iThemba LABS); SMIT, F.D. (iThemba LABS); STEYN, G.F. (iThemba LABS); USMAN, I. (University of the Witwatersrand); CARTER, J. (University of the Witwatersrand); VAN ZYL, J.J. (Stellenbosch University); BRUMMER, J.W. (Stellenbosch University); LI, K.C. (Stellenbosch University); DONALDSON, L. (iThemba LABS); FARBER, M. (University of Cologne); WIEDEKING, M. (iThemba LABS); WIENERT, M. (University of Cologne); ADSLEY, P. (University of the Witwatersrand and iThemba LABS); JONES, P. (iThemba LABS); PAPKA, P. (iThemba LABS); VON NEUMAN-COSEL, P. (Institut fuer Kernphysik, Technische Universitaet Darmstadt); MOLEMA, P.T. (University of the Witwatersrand); NEVELING, R. (iThemba LABS); JONGILE, S. (iThemba LABS); SIEM, S. (University of Oslo); TRIAMBAK, S. (University of the Western Cape); KHUMALO, T (iThemba LABS); PESUDO, V. (iThemba LABS)

Presenter: JIVAN, Harshna (University of the Witwatersrand)