Conference on Neutrino and Nuclear Physics (CNNP2020) Arabella Hotel and Spa, South Africa, 24-28 February 2020



Contribution ID: 67

Type: Oral

## Decay Characteristics of the Scissors Mode of 0vββ-Decay Partner Isotopes\*

Friday, 28 February 2020 09:00 (20 minutes)

The search for neutrinoless double beta  $(0\nu\beta\beta)$  decay, a process only allowed if the neutrino were a Majorana particle, recently gained much attention with numerous experiments being dedicated to its observation. It would demonstrate lepto-genesis in the universe and allow the determination of the neutrino mass from its decay rate. However, to quantitatively extract the neutrino mass or estimate decay rates a nuclear matrix element (NME) is required, which has to be calculated using nuclear structure models. One of them is the Interacting Boson Model 2 (IBM-2), which will be discussed below. Those calculations can be difficult because many of the 0vßβ-decay candidate nuclei lie in regions of the nuclear chart that feature shape coexistence, with the hypothesized 0vββ-decay mother nucleus <sup>150</sup>Nd and its daughter <sup>150</sup>Sm even being located in the region of a shape phase transition along their respective isotopic chains. In particular, the occurrence of shape coexistence may lead to a significant population of an excited  $0^+$  state in  $0\nu\beta\beta$  decay. To improve  $0\nu\beta\beta$ -NME calculations for <sup>150</sup>Nd and <sup>150</sup>Sm within the IBM-2 information on its so-called Majorana interaction is needed. Therefore, new data on the decay characteristics of the scissors mode of these nuclei was recently taken in nuclear resonance fluorescence experiments performed at the High Intensity y-ray Source. The decay characteristics of the scissors mode are sensitive to the nuclear deformation and allow inducing constraints on model parameters, especially the Majorana parameters of the IBM-2, in turn resulting in a more reliable prediction of the 0vββ-NME. Similar information has also been obtained for the 0vββ-partner nuclides <sup>82</sup>Se and <sup>82</sup>Kr. The experimental results and updated IBM-2 calculations will be presented and discussed. \*Supported by the DFG through the research grant SFB 1245 and by the State of Hesse under the grant "Nuclear Photonics" within the LOEWE program.

Primary author: KLEEMANN, Jörn (AG Pietralla Institut für Kernphysik Technische Universität Darmstadt)

**Co-authors:** BECK, T (IKP, TU Darmstadt, Germany); FRIMAN-GAYER, U (IKP, TU Darmstadt, Germany); PIETRALLA , N (IKP, TU Darmstadt, Germany); WERNER, V (IKP, TU Darmstadt, Germany); FINCH, S (TUNL, Durham NC, USA); KOTILA, J (University of Jyväskylä, Jyväskylä, Finland); FNU, Krishihayan (TUNL, Durham NC, USA); LÖHER, B (IKP, TU Darmstadt, Germany and GSI, Darmstadt, Germany); PAI, H (IKP, TU Darmstadt, Germany and SINP, Kolkata, India); PAPST, O (IKP, TU Darmstadt, Germany); TORNOW, W (TUNL, Durham NC, USA); WEINERT, M (IKP, Universität zu Köln, Germany)

Presenter: KLEEMANN, Jörn (AG Pietralla Institut für Kernphysik Technische Universität Darmstadt)

Session Classification: Contributed Talks